



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07K 7/08, 15/10, 15/28, C12N 15/29, 15/70, 37/02, A61K 37/02	A2	(11) International Publication Number: WO 94/21675 (43) International Publication Date: 29 September 1994 (29.09.94)
(21) International Application Number: PCT/US94/02537 (22) International Filing Date: 9 March 1994 (09.03.94) (30) Priority Data: 08/031,001 12 March 1993 (12.03.93) US (71) Applicant: IMMULOGIC PHARMACEUTICAL CORPORATION [US/US]; 610 Lincoln Street, Waltham, MA 02154 (US). (72) Inventors: GRIFFITH, Irwin, J.; 13 Southwick Road, North Reading, MA 01864 (US). KUO, Mei-chang; 5 Cox Road, Winchester, MA 01890 (US). LUQMAN, Mohammad; 13 Carriage Drive, Acton, MA 01720 (US). POWERS, Stephen, Palmer; 2008 Stearns Hill Road, Waltham, MA 02154 (US). (74) Agents: CHANNING, Stacey, L. et al.; Immulogic Pharmaceutical Corporation, 610 Lincoln Street, Waltham, MA 02154 (US).		(81) Designated States: AU, BB, BG, BR, BY, CA, CN, CZ, FL, HU, JP, KP, KR, KZ, LK, LV, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: T CELL EPITOPES OF RYEGRASS POLLEN ALLERGEN (57) Abstract <p>The present invention provides isolated peptides of <i>Lol p I</i>, a major protein allergen of the species <i>Lolium perenne</i>. Peptides within the scope of the invention comprise at least one T cell epitope, or preferably at least two T cell epitopes of a protein allergen of <i>Lol p I</i>. The invention also provides modified peptides having similar or enhanced therapeutic or diagnostic properties as the corresponding, naturally-occurring allergen or portion thereof, but having additional properties, e.g., reduced side effects. The invention further provides nucleic acid sequences coding for peptides of the invention. Methods of treatment and diagnosis of sensitivity to <i>Lol p I</i> or an allergen immunologically related to <i>Lol p I</i> in an individual (such as <i>Dac g I</i>, <i>Poa p I</i>, or <i>Phl p I</i>) also are provided. Compositions for therapeutic, diagnostic or reagent uses comprising one or more peptides of the invention are also provided.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

T CELL EPITOPES OF RYEGRASS POLLEN ALLERGEN

Background of the Invention

5 The most abundant proteins of grass pollen are allergens, which are the major cause of allergic disease in temperate climates (Marsh (1975), "Allergens and the genetics of allergy"; in M. Sela (ed), *The Antigens*, 3:271-359, Academic Press Inc., London, New York)., Hill *et al.* (1979) *Medical Journal of Australia*, 1:426-429). The first descriptions of the allergenic proteins in ryegrass showed that they are
10 immunochemically distinct, and are known as groups I, II, III and IV (Johnson and Marsh (1965), *Nature*, 206:935-942; and Johnson and Marsh (1966) *Immunochemistry*, 3:91-100). Using the International Union of Immunological Societies' (IUIS) nomenclature, these allergens are designated *Lol p* I, *Lol p* II, *Lol p* III, and *Lol p* IV. Another important *Lolium perenne* allergen which has been
15 identified in the literature is *Lol p* IX, also known as *Lol p* V or *Lol p* Ib, which has been found to be closely related to the Group V protein allergens in grasses.

 These proteins have been identified in pollen from ryegrass, *Lolium perenne*, and act as antigens in triggering immediate (Type 1) hypersensitivity in susceptible humans.

20 *Lol p* I is defined as an allergen because of its ability to bind to specific IgE in sera of ryegrass-sensitive patients, to act as an antigen in IgG responses and to trigger T-cell responses. The allergenic properties have been assessed by direct skin testing of grass pollen-sensitive patients. The results showed that 84% had a skin sensitivity to *Lol p* I (Freidhoff, et al., (1986) *J. Allergy Clin. Immunol.*, 78:1190-1201)
25 demonstrating the primary importance of this protein as the major allergen. Furthermore, 95% of patients demonstrated to be grass pollen-sensitive possessed specific IgE antibody that bound to *Lol p* I, as demonstrated by immunoblotting (Ford and Baldo (1986) *International Archives of Allergy and Applied Immunology*, 81:193-203).

30 Substantial allergenic cross-reactivity between grass pollens has been demonstrated using an IgE-binding assay, the radioallergo-sorbent test (RAST), for example, as described by Marsh *et al.* (1970) *J. Allergy*, 46:107-121, and Lowenstein (1978) *Prog. Allergy*, 25:1-62. (Karger, Basel).

 The immunochemical relationship of *Lol p* I with other grass pollen
35 antigens has been demonstrated using both polyclonal and monoclonal antibodies (e.g., Smart and Knox (1979) *International Archives of Allergy and*

Applied Immunology, **62**: 173-187; Singh and Knox (1985), *International Archives of Allergy and Applied Immunology*, **78**:300-304). Antibodies have been prepared to both purified proteins and IgE-binding components. These data demonstrate that the major allergen present in pollen of closely related
5 grasses is immunochemically similar to *Lol p I* (Singh and Knox, *supra*). Grasses that may be considered immunochemically related to *Lol p I* and that comprise allergens which may be considered immunologically cross-reactive with antibody to *Lol p I* include:

Pooid (festucoid) grasses of the Poaceae (Gramineae) family include the
10 following. GROUP 1: Triticaneae: *Bromus inermis*, smooth brome; *Agropyron repens*, English couch; *A. cristatum*; *Secale cereale* rye *Triticum aestivum*, wheat. GROUP 2: Poanae: *Dactylis glomerata*, orchard grass of cocksfoot; *Festuca elatior*, meadow fescue; *Lolium perenne*, perennial ryegrass; *L. multiflorum*, Italian ryegrass; *Poa pratensis*, Kentucky bluegrass;
15 *P. compressa*, flattened meadow grass; *Avena sativa*, oat; *Holcus lanatus*, velvet grass or Yorkshire fog; *Anthoxanthum odoratum*; sweet vernal grass; *Arrhenatherum elatius*, oat grass; *Agrostis alba*, red top; *Phleum pratense*, timothy; *Phalaris arundinacea*, reed canary grass. *Panicoid* grass, *Paspalum notatum*, Bahia grass, Andropogonoid grasses: *Sorghum halepensis*, Johnson
20 grass.

In view of the prevalence of ryegrass pollen allergens and related grass allergens all over the world, there is a pressing need for the development of compositions and methods that could be used in detecting sensitivities to *Lol p I* or other immunologically related grass allergens, or in treating sensitivities to
25 such allergens, or in assisting in the manufacture of medicaments to treat such sensitivities. The present invention provides materials and methods having one or more of those utilities.

Summary of the Invention

30 The present invention provides isolated peptides of *Lol p I*. Peptides within the scope of the invention comprise at least one T cell epitope, preferably at least two T cell epitopes of *Lol p I*. The invention further provides peptides comprising at least two regions, each region comprising at least one T cell epitope of *Lol p I*.

The invention also provides modified peptides having similar or enhanced therapeutic or diagnostic properties as the corresponding, naturally-occurring allergen or portion thereof, but also having advantageous physical or biological properties, such as reduced side effects, reduced IgE binding, improved solubility, increased *in vitro* or *in vivo* T cell stimulating ability, increased stability or the like. Preferred peptides of the invention are capable of modifying, in a *Lol p I*-sensitive individual to whom they are administered, the allergic response of the individual to *Lol p I* or an allergen immunologically cross-reactive with *Lol p I*, e.g., allergens derived from pollen belonging to the Poaceae (Gramineae) family, such as *Dactylis glomerata* (*Dac g I*), *Poa pretensis* (*Poa p I*) and *Phleum pratense* (*Phl p I*), as discussed above.

The present invention also provides non-native (i.e., recombinant or chemically synthesized) *Lol p I* peptides or their derivatives or homologues and provides non-native allergenic protein or peptides immunologically cross-reactive with antibodies or with T cells of *Lol p I* or derivatives or homologues thereof.

The present invention also provides *Dac g I* and *Poa p I* protein allergens which are immunologically cross-reactive with *Lol p I*, and fragments of *Dac g I* and *Poa p I* produced in a host cell transformed with a nucleic acid sequence coding for *Dac g I* and *Poa p I*, respectively, and fragments of *Dac g I* and *Poa p I* prepared synthetically. The present invention further provides nucleic acid sequences coding for *Dac g I*, *Poa p I* and fragments thereof. Also provided are isolated peptides of *Dac g I* and *Poa p I* comprising at least one T cell epitope which are immunologically cross-reactive with peptides comprising at least one T cell epitope derived from *Lol p I*.

Methods of treatment and of diagnosis of sensitivity to ryegrass pollen protein, *Lol p I*, or to pollen proteins that are immunologically related to *Lol p I* (such as *Dac g I*, *Phl p I* and *Poa p I*), as well as compositions comprising one or more peptides of the invention, are also provided.

Further features of the present invention will be better understood from the following detailed description of the preferred embodiments of the invention in conjunction with the appended figures.

Brief Description of the Figures

Fig. 1 shows the nucleotide sequence of cDNA clone 26.j (SEQ ID NO 1) and its predicted amino acid sequence (SEQ ID NO: 2). Clone 26.j is a PCR-generated, full-length clone of *Lol p I*.

Fig. 2 shows various peptides of desired lengths derived from *Lol p I* (SEQ ID NO: 3-30); such peptides include polymorphisms inherent in the *Lol p I* sequence (i.e., LPI-4.1 (SEQ ID NO: 8) and LPI-16.1 (SEQ ID NO: 23)) or homologues of peptides derived from *Lol p I* (i.e., LPI-11 (SEQ ID NO: 15), and LPI-12 (SEQ ID NO: 17)).

Fig. 3 is a graphic representation depicting responses of T cell lines from thirty-five grass-sensitive patients primed *in vitro* with purified native *Lol p I* and analyzed for response to various *Lol p I* peptides by percent of positive responses (with an S.I. of at least two, shown over each bar), the mean stimulation index of positive response for the peptide (shown over each bar in parentheses) and the positivity index (% positive x mean S.I. index, Y axis).

Fig. 4 shows various peptides of desired lengths derived from *Lol p I* (SEQ ID NO: 23, 25, 27, 30-50).

Fig. 5 shows the nucleotide sequence of cDNA clone 106.5 (SEQ ID NO: 51) and its predicted amino acid sequence (SEQ ID NO: 52). Clone 106.5 is a PCR-generated, full-length clone of *Dac g I*.

Fig. 6 shows the nucleotide sequence of cDNA clone 114 (SEQ ID NO: 53) and its predicted amino acid sequence (SEQ ID NO: 54). Clone 114 is a PCR-generated, full-length clone of *Poa p I*.

Fig. 7 shows the nucleotide sequence of cDNA clone 20 (SEQ ID NO: 55) and its predicted amino acid sequence (SEQ ID NO: 56). Clone 20 is a PCR generated, full length clone of *Phl p I*.

Fig. 8 shows a comparison of the amino acid sequences of the mature protein of *Lol p I* (SEQ ID NO: 57), *Dac g I* (SEQ ID NO: 58), *Phl p I* (SEQ ID NO: 59), and *Poa p I* (SEQ ID NO: 60), including polymorphisms thereof.

Fig. 9 shows a comparison of various peptides comprising at least one T cell epitope derived from *Lol p I*, with homologous peptides derived from the same regions of *Dac g I*, *Phl p I*, and *Poa p I* (SEQ ID NO: 23, 25, 27, 30, 61-70).

Detailed Description of the Invention

The present invention provides isolated peptides derived from *Lol p I* (SEQ ID NO: 3-50). The present invention also provides *Dac g I* and *Poa p I* protein allergens which are immunologically cross-reactive with *Lol p I*. The term "peptide"

as used herein refers to any protein fragment of *Lol p I* that induces an immune response. The terms "fragment" and "antigenic fragment" of a protein as used interchangeably herein refer to an amino acid sequence having fewer amino acid residues than the entire native amino acid sequence of the protein from which the
5 fragment is derived, and that induces an immune response. The terms "isolated" and "purified" as used herein refer to peptides of the invention which are substantially free of cellular material or culture medium when produced by recombinant DNA techniques, or substantially free of chemical precursors or other chemicals when synthesized chemically. Preferred peptides of the invention include peptides derived
10 from *Lol p I* which comprise at least one T cell epitope of the allergen, or a portion of such a peptide which includes at least one T cell epitope.

Peptides comprising at least two regions, each region comprising at least one T cell epitope *Lol p I* are also within the scope of the invention. Isolated peptides or regions of isolated peptides, each comprising at least two T cell epitopes of the *Lol p*
15 *I* protein allergen are particularly desirable for increased therapeutic effectiveness. Peptides that are immunologically related (e.g., by antibody or T cell cross-reactivity) to peptides of the present invention, such as peptides derived from *Dac g I* and *Poa p I*, are also within the scope of the invention. Peptides immunologically related by antibody cross-reactivity are recognized by antibodies specific for a peptide of *Lol p I*.
20 Peptides immunologically related to a given peptide by T cell cross-reactivity are capable of also reacting with the same T cells that react with that given peptide.

Isolated protein and peptides of the invention can be produced by recombinant DNA techniques in a host cell transformed with a nucleic acid having a sequence encoding such peptide. The isolated peptides of the invention can also be produced by
25 chemical synthesis. When a protein or peptide is produced by recombinant techniques, host cells transformed with a nucleic acid having a sequence encoding a peptide of the invention or the functional equivalent of the nucleic acid sequence are cultured in a medium suitable for the cells. Peptides can be purified from cell culture medium, host cells, or both, using techniques known in the art for purifying peptides
30 and proteins including ion-exchange chromatography, gel filtration chromatography, ultrafiltration, electrophoresis or immunopurification with antibodies specific for the peptide, the protein allergen from which the peptide is derived, or a portion thereof.

The present invention provides expression vectors and host cells transformed to express the nucleic acid sequences of the invention. Nucleic acids coding for *Lol p I*
35 peptides of the invention, or at least a portion thereof, may be expressed in bacterial

cells such as *E. coli*, insect cells, yeast, or mammalian cells such as Chinese hamster ovary cells (CHO). Suitable expression vectors, promoters, enhancers, and other expression control elements may be found in Sambrook et al. *Molecular Cloning: A Laboratory Manual*, second edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989. Other suitable expression vectors, promoters, enhancers, and other expression elements are known to those skilled in the art. Expression in mammalian, yeast or insect cells leads to partial or complete glycosylation of the recombinant material and formation of any inter- or intra-chain disulfide bonds. Suitable vectors for expression in yeast include YepSec1 (Baldari et al. (1987) *Embo J.*, **6**: 229-234); pMFa (Kurjan and Herskowitz (1982) *Cell*, **30**: 933-943); JRY88 (Schultz et al. (1987) *Gene*, **54**: 113-123) and pYES2 (Invitrogen Corporation, San Diego, CA). These vectors are freely available. Baculovirus and mammalian expression systems are also available. For example, a baculovirus system is commercially available (PharMingen, San Diego, CA) for expression in insect cells while the pMSG vector is commercially available (Pharmacia, Piscataway, NJ) for expression in mammalian cells.

For expression in *E. coli*, suitable expression vectors include, among others, pTRC (Amann et al. (1988) *Gene*, **69**: 301-315); pGEX (Amrad Corp., Melbourne, Australia); pMAL (N.E. Biolabs, Beverly, MA); pRIT5 (Pharmacia, Piscataway, NJ); pET-11d (Novagen, Madison, WI) Jameel et al., (1990) *J. Virol.*, **64**:3963-3966; and pSEM (Knapp et al. (1990) *BioTechniques*, **8**: 280-281). The use of pTRC, and pET-11d, for example, will lead to the expression of unfused protein. The use of pMAL, pRIT5 pSEM and pGEX will lead to the expression of allergen fused to maltose E binding protein (pMAL), protein A (pRIT5), truncated β -galactosidase (PSEM), or glutathione S-transferase (pGEX). When a *Lol p I* peptide of the invention, is expressed as a fusion protein, it is particularly advantageous to introduce an enzymatic cleavage site at the fusion junction between the carrier protein and the *Lol p I* peptide. The *Lol p I* peptide may then be recovered from the fusion protein through enzymatic cleavage at the enzymatic site and biochemical purification using conventional techniques for purification of proteins and peptides. Suitable enzymatic cleavage sites include those for blood clotting Factor Xa or thrombin for which the appropriate enzymes and protocols for cleavage are commercially available from, for example, Sigma Chemical Company, St. Louis, MO and N.E. Biolabs, Beverly, MA. The different vectors also have different promoter regions allowing constitutive or inducible expression with, for example, IPTG induction (PRTC, Amann et al., (1988)

supra; pET-11d, Novagen, Madison, WI) or temperature induction (pRIT5, Pharmacia, Piscataway, NJ). It may also be appropriate to express recombinant *Lol p* I peptides in different *E. coli* hosts that have an altered capacity to degrade recombinantly expressed proteins (e.g., U.S. Patent 4,758,512). Alternatively, it may be advantageous to alter the nucleic acid sequence to use codons preferentially utilized by *E. coli*, where such nucleic acid alteration would not affect the amino acid sequence of the expressed protein.

Host cells can be transformed to express the nucleic acid sequences of the invention using conventional techniques such as calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, or electroporation. Suitable methods for transforming the host cells may be found in Sambrook et al. *supra*, and other laboratory textbooks. The nucleic acid sequences of the invention may also be chemically synthesized using standard techniques (i.e., solid phase synthesis). Details of the cloning of *Lol p* I are given in the Examples.

Inducible non-fusion expression vectors include pTrc (Amann *et al.*, (1988) *Gene*, **69**:301-315) and pET11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology*, Academic Press, San Diego, California (1990), **185**:60-89). While target gene expression relies on host RNA polymerase transcription from the hybrid trp-lac fusion promoter in pTrc, expression of target genes inserted into pET11d relies on transcription from the T7 *gn10-lac 0* fusion promoter mediated by coexpressed viral RNA polymerase (T7 *gn1*). This viral polymerase is supplied by host strains BL21(DE3) or HMS174(DE3) from a resident λ prophage harboring a T7 *gn1* under the transcriptional control of the lacUV 5 promoter.

One strategy to maximize recombinant *Lol p* I peptide expression in *E. coli* is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein (Gottesman, S., *Gene Expression Technology: Methods in Enzymology*, Academic Press, San Diego, California (1990), **185**:119-128). Another strategy would be to alter the nucleic acid sequence of the desired gene to be inserted into an expression vector so that the individual codons for each amino acid would be those preferentially utilized in highly expressed *E. coli* proteins (Wada *et al.*, (1992) *Nuc. Acids Res.*, **20**:2111-2118). Such alteration of nucleic acid sequences of the invention could be carried out by standard DNA synthesis techniques.

The nucleic acids of the invention can also be chemically synthesized using standard techniques. Various methods of chemically synthesizing

polydeoxynucleotides are known, including solid-phase synthesis which, like peptide synthesis, has been fully automated in commercially available DNA synthesizers (See e.g., Itakura *et al.* U.S. Patent 4,598,049; Caruthers *et al.* U.S. Patent 4,458,066; and Itakura U.S. Patents 4,401,796 and 4,373,071, incorporated by reference herein).

5 The present invention also provides fragments of nucleic acid sequences encoding peptides of the invention. As used herein, the term "fragment" of a nucleic acid sequence refers to a nucleotide sequence having fewer bases than the nucleotide sequence coding for the entire amino acid sequence of the protein. Nucleic acid sequences used in any embodiment of this invention can be cDNA obtained as
10 described herein, or alternatively, can be any oligodeoxynucleotide sequence having all or a portion of a sequence represented herein, or their functional equivalents. Such oligodeoxynucleotide sequences can be produced chemically or mechanically, using known techniques. A functional equivalent of an oligonucleotide sequence of *Lol p I* is one which is 1) a sequence capable of hybridizing to a complementary
15 oligonucleotide to which the sequence (or corresponding sequence portions) of *Lol p I* as shown in Fig. 1 (SEQ ID NO: 1) or fragments thereof hybridizes, or 2) the sequence (or corresponding sequence portion) complementary to the sequence of *Lol p I* as shown in Fig. 1 (SEQ ID NO: 1), and/or 3) a sequence which encodes a product (e.g., a polypeptide or peptide) having the same functional characteristics of the
20 product encoded by the sequence (or corresponding sequence portion) of *Lol p I* as shown in Fig. 1 (SEQ ID NO: 1). Whether a functional equivalent must meet one or both criteria will depend on its use (e.g., if it is to be used only as an oligonucleotide probe, it need meet only the first or second criteria and if it is to be used to produce a *Lol p I* peptide of the invention, it need only meet the third criterion).

25 Preferred nucleic acids encode a peptide having at least about 50% homology to a *Lol p I* peptide of the invention, more preferably at least about 60% homology and most preferably at least about 70% homology with a *Lol p I* peptide of the invention. Nucleic acids that encode peptides having at least about 90%, more preferably at least about 95%, and most preferably at least about 98-99% homology with *Lol p I* peptides
30 of the invention are also within the scope of the invention. Homology refers to sequence similarity between two peptides of *Lol p I*, or between two nucleic acid molecules. Homology can be determined by comparing a position in each sequence which may be aligned for purposes of comparison. When a position in the compared sequence is occupied by the same nucleotide or amino acid, then molecules are

homologous at that position. A degree of homology between sequences is a function of the number of matching or homologous positions shared by the sequences.

Preferred nucleic acid fragments encode peptides of at least 7 amino acid residues in length, and preferably 13-40 amino acid residues in length, and more preferably at least 16-30 amino acid residues in length. Nucleic acid fragments encoding peptides of at least 30 amino acid residues in length, at least 40 amino acid residues in length, at least about 80 amino acid residues in length, at least about 100 amino acid residues in length or more, are also contemplated.

Also within the scope of the invention are nucleic acid sequences encoding allergens immunologically cross-reactive with *Lol p I*, such as full length *Dac g I* and *Poa p I* proteins or peptides (Figs 5 (SEQ ID NO: 52), 6 (SEQ ID NO: 54), and 9 (SEQ ID NO: 23, 25, 27, 30, 61-70)). Proteins and peptides of *Dac g I* and *Poa p I* may be produced recombinantly as discussed above, or synthetically. Expression vectors and host cells transformed to express *Dac g I* and *Poa p I* proteins or peptides thereof are also within the scope of the invention. Details of the cloning of *Dac g I* and *Poa p I* are given in the examples.

The present invention also provides a method of producing isolated *Lol p I* peptides of the invention or a portion thereof, comprising the steps of culturing a host cell transformed with a nucleic acid sequence encoding a *Lol p I* peptide of the invention in an appropriate medium to produce a mixture of cells and medium containing said *Lol p I* peptide; and purifying the mixture to produce substantially pure *Lol p I* peptide. Host cells transformed with an expression vector containing DNA coding for a *Lol p I* peptide of the invention are cultured in a suitable medium for the host cell. *Lol p I* peptides of the invention can be purified from cell culture medium, host cells, or both using techniques known in the art for purifying peptides and proteins including ion-exchange chromatography, gel filtration chromatography, ultrafiltration, electrophoresis and immunopurification with antibodies specific for the *Lol p I* peptides or portions thereof.

Another aspect of the present invention pertains to an antibody specifically reactive with a *Lol p I* peptide. Such antibodies may be used to standardize allergen extracts or to isolate the naturally occurring *Lol p I*. Also, *Lol p I* peptides of the invention can be used as "purified" allergens to standardize allergen extracts. For example, an animal such as a mouse or rabbit can be immunized with an immunogenic form of an isolated *Lol p I* peptide of the invention capable of eliciting an antibody response. Techniques for conferring immunogenicity on a peptide include conjugation

to carriers or other techniques well-known in the art. The *Lol p I* peptide can be administered in the presence of adjuvant. The progress of immunization can be monitored by detection of antibody titers in plasma or serum standard ELISA or other immunoassay can be used with the immunogen as antigen to assess the levels of antibodies.

Following immunization, anti-*Lol p I* peptide antisera can be obtained and, if desired, polyclonal anti-*Lol p I* peptide antibodies from the serum. To produce monoclonal antibodies, antibody producing cells (lymphocytes) can be harvested from an immunized animal and fused by standard somatic cell fusion procedures with immortalizing cells such as myeloma cells to yield hybridoma cells. Hybridoma cells can be screened immunochemically for production of antibodies reactive with the *Lol p I* peptides of the invention. These sera or monoclonal antibodies can be used to standardize allergen extracts.

Through use of the peptides and antibodies of the present invention, preparations of consistent, well-defined composition and uniform biological activity can be made. Compositions having therapeutic activity may be administered for therapeutic purposes (e.g., to modify the allergic response of a ryegrass sensitive individual to pollen of such grasses or pollen of an immunologically related grass such as *Dac g I*, *Poa p I* and *Phl p I*). Administration of such peptides may, for example, modify B-cell response to *Lol p I* allergen, T-cell response to *Lol p I* allergen or both responses. Isolated peptides can also be used to study the mechanism of immunotherapy of ryegrass pollen allergy and to design modified derivatives or analogues useful in immunotherapy. Compositions according to the invention will have utility in diagnosis of ryegrass sensitivity or sensitivity to grass allergens cross-reactive to ryegrass allergens, because the components include T cell epitopes recognizing the allergens.

The present invention also pertains to T cell clones which specifically recognize *Lol p I* peptides of the invention. These T cell clones may be suitable for isolation and molecular cloning of the gene for the T cell receptor which is specifically reactive with a peptide of the present invention. The T cell clones may be produced as described in Example 4, or as described in *Cellular Molecular Immunology*, Abdul K. Abbas et al., W.B. Saunders Co. (1991) pg. 139. The present invention also pertains to soluble T cell receptors. These receptors may inhibit antigen-dependent activation of the relevant T cell subpopulation within an individual sensitive to *Lol p I*. Antibodies specifically reactive with such a T cell receptor can also be produced according to the

techniques described herein. Such antibodies may also be useful to block T-cell-MHC interaction in an individual. Methods for producing soluble T cell receptors are described in *Immunology: A Synthesis*, 2nd Ed., Edward S. Golub *et al.*, Sinaur Assoc., Sunderland, Massachusetts, (1991) pp. 366-369.

5 It is also possible to modify the structure of a peptide of the invention to achieve additional advantageous physical or biological properties such as increasing solubility, enhancing therapeutic or preventive efficacy, increasing stability (e.g., shelf life *ex vivo* or resistance to proteolytic degradation *in vivo*), decreasing adverse side effects, and the like. A modified peptide can be produced in which the amino acid
10 sequence has been altered, such as by amino acid substitution, deletion, or addition, in order to modify immunogenicity and/or to reduce allergenicity. Peptides may also be advantageously modified by addition or conjugation with another peptide or other component.

 For example, a peptide can be modified so that it maintains the ability to induce
15 T cell anergy and to bind MHC proteins but reduces the ability to induce a strong proliferative response, or possibly any proliferative response, when administered in immunogenic form. In this instance, critical binding residues for the T cell receptor can be determined using known techniques (e.g., substitution of each residue and determination of the presence or absence of T cell reactivity). Those residues shown
20 to be essential to interact with the T cell receptor can be modified by replacing the essential amino acid with another preferably similar amino acid residue (a "conservative substitution") whose presence is shown to enhance, diminish but not eliminate, or not affect T cell reactivity. In addition, those amino acid residues that are not essential for T cell receptor interaction can be modified by replacement with
25 another amino acid whose incorporation may enhance, diminish or not affect T cell reactivity but does not eliminate binding to relevant MHC.

 Additionally, peptides of the invention can be modified by replacing an amino acid shown to be essential to interact with the MHC protein complex with another, preferably similar amino acid residue (conservative substitution) whose presence is
30 shown to enhance, diminish but not eliminate or not affect T cell reactivity. In addition, amino acid residues that are not essential for interaction with the MHC protein complex but that still bind the MHC protein complex can be modified by replacement with another amino acid whose incorporation may enhance, not affect, or diminish but not eliminate T cell reactivity. Preferred amino acid substitutions for non-

essential amino acids include, but are not limited to substitutions with alanine, glutamic acid, or a methyl amino acid.

In order to enhance stability and/or reactivity, peptides of the invention can also be modified to incorporate one or more polymorphisms in the amino acid
5 sequence of the protein allergen resulting from natural allelic variation. Additionally, D-amino acids, non-natural amino acids or non-amino acid analogues can be substituted or added to produce a modified peptide within the scope of this invention. Furthermore, peptides of the present invention can be modified using the polyethylene glycol (PEG) method of A. Schon and co-workers (Wie et al., *supra*) to produce a
10 protein or peptide conjugated with PEG. In addition, PEG can be added during chemical synthesis of a protein or peptide of the invention. Modifications of peptides or portions thereof can also include reduction/ alyklation (Tarr in: *Methods of Protein Microcharacterization*, J.E. Silver ed. Humana Press, Clifton, NJ, pp 155-194 (1986)); acylation (Tarr, *supra*); chemical coupling to an appropriate carrier (Mishell and Shiigi,
15 eds, *Selected Methods in Cellular Immunology*, WH Freeman, San Francisco, CA (1980); U.S. Patent 4,939,239; or mild formalin treatment (Marsh *International Archives of Allergy and Applied Immunology*, **41**:199-215 (1971)).

To facilitate purification and potentially increase solubility of peptides of the invention, it is possible to add reporter group(s) to the peptide backbone. For
20 example, poly-histidine can be added to a peptide to purify the peptide by immobilized metal ion affinity chromatography (Hochuli, E. et al., *Bio/Technology*, **6**:1321-1325 (1988)). In addition, specific endoprotease cleavage sites can be introduced, if desired, between a reporter group and amino acid sequences of a peptide to facilitate isolation of peptides free of irrelevant sequences. In order to successfully desensitize
25 an individual to a protein antigen, it may be necessary to increase the solubility of a peptide by adding functional groups to the peptide or by not including hydrophobic T cell epitopes or regions containing hydrophobic epitopes in the peptides or hydrophobic regions of the protein or peptide. Functional groups such as charged amino acid pairs (e.g., KK or RR) are particularly useful for increasing the solubility
30 of a peptide when added to the amino or carboxy terminus of the peptide. Examples of modifications to peptides to increase solubility include modifications to peptide LPI-16.1 (SEQ ID NO: 23) (Fig. 2), such modified peptides include: LPI-16.2 (SEQ ID NO: 31), LPI-16.3 (SEQ ID NO: 32), LPI-16.4 (SEQ ID NO 33) , LPI-16.5 (SEQ ID NO: 34), LPI-16.6 (SEQ ID NO: 35), LPI-16.7 (SEQ ID NO: 36) , LPI-16.9
35 (SEQ ID NO: 37), LPI-16.10 (SEQ ID NO: 38), all as shown in Fig. 4.

To potentially aid proper antigen processing of T cell epitopes within a peptide, canonical protease sensitive sites can be recombinantly or synthetically engineered between regions, each comprising at least one T cell epitope. For example, charged amino acid pairs, such as KK or RR, can be introduced between regions within a peptide during recombinant construction of the peptide or added to the amino or carboxy terminus of a synthetically produced peptide. The resulting peptide can be rendered sensitive to cathepsin and/or other trypsin-like enzymes cleavage to generate portions of the peptide containing one or more T cell epitopes. In addition, as mentioned above, such charged amino acid residues can result in an increase in solubility of a peptide.

Site-directed mutagenesis of DNA encoding a peptide of the invention can be used to modify the structure of the peptide by methods known in the art. Such methods may, among others, include PCR with degenerate oligonucleotides (Ho et al., *Gene*, **77**:51-59 (1989)) or total synthesis of mutated genes (Hostomsky, Z. et al., *Biochem. Biophys. Res. Comm.*, **161**:1056-1063 (1989)). To enhance bacterial expression, the aforementioned methods can be used in conjunction with other procedures to change the eucaryotic codons in DNA constructs encoding protein or peptides of the invention to ones preferentially processed in *E. coli*, yeast, mammalian cells, or other prokaryotic or eukaryotic host cells.

Peptides of the present invention can also be used for detecting and diagnosing ryegrass pollinosis. For example, this could be done *in vitro* by combining blood or blood products obtained from an individual to be assessed for sensitivity to ryegrass pollen or another cross-reactive pollen such as *Dac g* I, *Poa p* I and *Phl p* I, with an isolated peptide(s) of *Lol p* I, under conditions appropriate for binding of components in the blood (e.g., antibodies, T-cells, B cells) with the peptide(s) and determining the extent to which such binding occurs. Other diagnostic methods for allergic diseases in which the protein, peptides or antibodies of the present invention will be useful include radio-allergosorbent test (RAST), paper radioimmunosorbent test (PRIST), enzyme linked immunosorbent assay (ELISA), radioimmunoassays (RIA), immuno-radiometric assays (IRMA), luminescence immunoassays (LIA), histamine release assays and IgE immunoblots.

The presence in individuals of IgE specific for at least one protein allergen and the ability of T cells of the individuals to respond to T cell epitope(s) of the protein allergen can be determined by administering to the individuals an Immediate Type Hypersensitivity test and a Delayed Type Hypersensitivity test. The individuals

are administered an Immediate Type Hypersensitivity test (see e.g., *Immunology* (1985) Roitt, I.M., Brostoff, J., Male, D.K. (eds), C.V. Mosby Co., Gower Medical Publishing, London, NY, pp. 19.2-19.18; pp. 22.1-22.10) utilizing the protein allergen or a portion thereof, or a modified form of the protein allergen or a portion thereof, each of which binds IgE specific for the allergen. The same individuals are administered a Delayed Type Hypersensitivity test prior to, simultaneously with, or subsequent to administration of the Immediate Type Hypersensitivity test. Of course, if the Immediate Type Hypersensitivity test is administered prior to the Delayed Type Hypersensitivity test, the Delayed Type Hypersensitivity test would be given to those individuals exhibiting a specific Immediate Type Hypersensitivity reaction. The Delayed Type Hypersensitivity test utilizes a modified form of the protein allergen or a portion thereof, the protein allergen produced recombinantly, or a peptide derived from the protein allergen, each of which has human T cell stimulating activity and each of which does not bind IgE specific for the allergen in a substantial percentage of the population of individuals sensitive to the allergen (e.g., at least about 75%). Those individuals found to have both a specific Immediate Type Hypersensitivity reaction and a specific Delayed Type Hypersensitivity reaction may be treated with a therapeutic composition comprising the same modified form of the protein or portion thereof, the recombinantly produced protein allergen, or the peptide, each as used in the Delayed Type Hypersensitivity test.

Isolated peptides of the invention, when administered in a therapeutic regimen to a *Lol p I*-sensitive individual (or an individual allergic to an allergen cross-reactive with ryegrass pollen allergen such as *Dac g I*, *Poa p I* and *Phl p I*) are capable of modifying the allergic response of the individual to *Lol p I* ryegrass pollen allergen (or such cross-reactive allergen). Preferably peptides of this invention are capable of modifying the B-cell response, T-cell response or both the B-cell and the T-cell response of the individual to the allergen. As used herein, modification of the allergic response of an individual sensitive to a ryegrass pollen allergen or cross-reactive allergen can be defined as non-responsiveness or diminution in symptoms to the allergen, as determined by standard clinical procedures (See, e.g., Varney et al, *British Medical Journal*, **302**:265-269 (1990)) including diminution in ryegrass pollen-induced asthmatic symptoms. As referred to herein, a diminution in symptoms includes any reduction in allergic response of an individual to the allergen after the individual has completed a treatment regimen with a peptide or protein of the invention. This diminution may be subjective (i.e., the patient feels more comfortable

in the presence of the allergen), or diminution in symptoms may be determined clinically, using standard skin tests known in the art and discussed above.

Lol p I peptides of the present invention having T cell stimulating activity, and thus comprising at least one T cell epitope, are particularly preferred. In referring to an epitope, the epitope will be the basic element or smallest unit of recognition by a receptor, particularly immunoglobulins, histocompatibility antigens and T cell receptors where the epitope comprises amino acids essential to receptor recognition. Amino acid sequences which mimic those of the epitopes and which are capable of down-regulating or reducing allergic response to *Lol p I* can also be used. T cell epitopes are believed to be involved in initiation and perpetuation of the immune response to a protein allergen that is responsible for the clinical symptoms of allergy. Such T cell epitopes are thought to trigger early events at the level of the T helper cell by binding to an appropriate HLA molecule on the surface of an antigen presenting cell and stimulating the relevant T cell subpopulation. These events lead to T cell proliferation, lymphokine secretion, local inflammatory reactions, recruitment of additional immune cells to the site, and activation of the B cell cascade leading to production of antibodies. One isotype of these antibodies, IgE, is fundamentally important to the development of allergic symptoms, and its production is influenced early in the cascade of events, at the level of the T helper cell, by the nature of the lymphokines secreted.

Exposure of ryegrass pollen-sensitive patients or patients sensitive to an immunologically cross-reactive protein allergen such as *Dac g I*, *Poa p I* and *Phl p I*, to isolated *Lol p I* peptides of the present invention which comprise at least one T cell epitope and are derived from *Lol p I* protein allergen, may tolerize or anergize appropriate T cell subpopulations such that they become unresponsive to the protein allergen and do not participate in stimulating an immune response upon such exposure. In addition, administration of a peptide of the invention or portion thereof which comprises at least one T cell epitope may modify the lymphokine secretion profile as compared with exposure to the naturally-occurring *Lol p I* protein allergen or portion thereof (e.g., may result in a decrease of IL-4 and/or an increase in IL-2). Furthermore, exposure to such peptide of the invention may influence T cell subpopulations which normally participate in the response to the naturally occurring allergen such that these T cells are drawn away from the site(s) of normal exposure to the allergen (e.g., nasal mucosa, skin, and lung) towards the site(s) of therapeutic administration of the fragment or protein allergen. This redistribution of T cell

subpopulations can have the effect of ameliorating or reducing the ability of an individual's immune system to stimulate the usual immune response at the site of normal exposure to the allergen, resulting in a diminution in allergic symptoms.

The isolated *Lol p I* peptides of the invention can be used in methods of
5 diagnosing, treating or preventing allergic reactions to *Lol p I* allergen or an immunologically related protein allergen such as *Dac g I*, *Poa p I* and *Phl p I*. Thus, the present invention provides compositions useful in allergy diagnosis and/or useful in allergy therapy comprising isolated *Lol p I* peptides or portions thereof. Such compositions will typically also comprise a pharmaceutically acceptable carrier or
10 diluent when intended for *in vivo* administration. Therapeutic compositions of the invention may include synthetically prepared *Lol p I* peptides.

Administration of the therapeutic compositions of the present invention to an individual to be desensitized can be carried out using known techniques. *Lol p I* peptides or portions thereof may be administered to an individual in combination with,
15 for example, an appropriate diluent, a carrier and/or an adjuvant. Pharmaceutically acceptable diluents include saline and aqueous buffer solutions. Pharmaceutically acceptable carriers include polyethylene glycol (Wie *et al.* (1981) *Int. Arch. Allergy Appl. Immunol.*, **64**:84-99) and liposomes (Strejan *et al.* (1984) *J. Neuroimmunol.*, **7**:27). For purposes of inducing T cell anergy, the therapeutic composition is
20 preferably administered in nonimmunogenic form, i.e., it does not contain adjuvant. The therapeutic compositions of the invention are administered to ryegrass pollen-sensitive individuals or individuals sensitive to an allergen which is immunologically cross-reactive with ryegrass pollen allergen (i.e., *Dactylis glomerata*, or *Sorghum halepensis*, etc.). Therapeutic compositions of the invention may also be used in the
25 manufacture of medicaments for treating sensitivity to ryegrass pollen allergen or an immunologically related pollen allergen.

Administration of the therapeutic compositions of the present invention to an individual to be desensitized can be carried out using known procedures at dosages and for periods of time effective to reduce sensitivity (i.e., to reduce the allergic
30 response) of the individual to the allergen. Effective amounts of the therapeutic compositions will vary according to factors such as the degree of sensitivity of the individual to ryegrass pollen, the age, sex, and weight of the individual, and the ability of the protein or fragment thereof to elicit an antigenic response in the individual.

The active compound (i.e., protein or fragment thereof) may be administered
35 in any convenient manner such as by injection (subcutaneous, intravenous, etc.), oral

administration, inhalation, transdermal application, or rectal administration.

Depending on the route of administration, the active compound may be coated within a material to protect the compound from the action of enzymes, acids and other natural conditions which may inactivate the compound.

5 For example, preferably about 1 µg- 3 mg and more preferably from about 20-750 µg of active compound (i.e., protein or fragment thereof) per dosage unit may be administered by injection. Dosage regimen may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the
10 therapeutic situation.

 To administer a peptide by other than parenteral administration, it may be necessary to coat the protein with, or co-administer the protein with, a material to prevent its inactivation. For example, the peptide or portion thereof may be co-administered with enzyme inhibitors or in liposomes. Enzyme inhibitors include
15 pancreatic trypsin inhibitor, diisopropylfluorophosphate (DEP) and trasylol. Liposomes include water-in-oil-in-water CGF emulsions as well as conventional liposomes (Strejan *et al.*, (1984), *J. Neuroimmunol.*, 7:27).

 The active compound may also be administered parenterally or intraperitoneally. Dispersions can also be prepared in glycerol, liquid polyethylene
20 glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations may contain a preservative to prevent the growth of microorganisms.

 Pharmaceutical compositions suitable for injection include sterile aqueous solutions (where the peptides are water soluble) or dispersions and sterile powders for
25 the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases, the composition intended for *in vivo* use must be sterile and must be fluid to the extent necessary to provide easy syringability. It should preferably be stable under the conditions of manufacture and storage and be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or
30 dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion, and by the use of surfactants. Prevention of the
35 action of microorganisms can be achieved by various antibacterial and antifungal

agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol and sorbitol or sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought
5 about, including in the composition, an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound (i.e., protein or peptide) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by
10 filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active
15 ingredient (i.e., protein or peptide) plus any additional desired ingredient from a previously sterile-filtered solution thereof.

When a peptide of the invention is suitably protected, as described above, the peptide may be orally administered, for example, with an inert diluent or an assimilable edible carrier. The peptide and other ingredients may also be enclosed in a hard or soft
20 gelatin capsule, compressed into tablets, or incorporated directly into the individual's food. For oral therapeutic administration, the active compound may be formulated with conventional excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 1% by weight of active compound. The
25 percentage of the composition and preparations may, of course, be varied and may conveniently be between about 5 to 80% by weight of the dosage unit. The amount of active compound in such therapeutically useful compositions is such that a suitable dosage will be obtained. Preferred compositions or preparations according to the present invention are prepared so that an oral dosage unit contains from about 10 μ g
30 to about 200 mg of active compound.

The tablets, troches, pills, capsules and the like may also contain the following: a binder such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such
35 as sucrose, lactose or saccharin or a flavoring agent such as peppermint, oil of

wintergreen, or cherry flavoring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar
5 or both. A syrup or elixir may contain the active compound, sucrose as a sweetening agent, methyl and propylparabens as preservative, a dye and flavoring such as cherry or orange flavor. Of course, any material used in preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compound may be incorporated into sustained-release
10 preparations and formulations.

As used herein "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any
15 conventional media or agent is incompatible with the active compound, use thereof in the therapeutic compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

Various isolated peptides of the invention derived from ryegrass pollen protein *Lol p I* are shown in Figs. 2 and 4 (SEQ ID NO: 3-50). Peptides comprising at least
20 two regions, each region comprising at least one T cell epitope of *Lol p I* are also within the scope of the invention. As used herein a region may include the amino acid sequence of a peptide of the invention as shown in Figs. 2 and 4 (SEQ ID NO: 3-50) or the amino acid sequence of a portion of such peptide.

To obtain isolated peptides of the present invention, *Lol p I* is divided into non-
25 overlapping peptides of desired length or overlapping peptides of desired lengths as discussed in Example 4 which can be produced recombinantly, or synthetically. Peptides comprising at least one T cell epitope are capable of eliciting a T cell response, such as T cell proliferation or lymphokine secretion, and/or are capable of inducing T cell anergy (i.e., tolerization). To determine peptides comprising at least
30 one T cell epitope, isolated peptides are tested by, for example, T cell biology techniques, to determine whether the peptides elicit a T cell response or induce T cell anergy. Those peptides found to elicit a T cell response or to induce T cell anergy are defined as having T cell stimulating activity.

As discussed in Example 4, human T cell stimulating activity can be tested by
35 culturing T cells obtained from an individual sensitive to *Lol p I* allergen, (i.e., an

individual who has an IgE-mediated immune response to *Lol p I* allergen) with a peptide derived from the allergen, then determining whether proliferation of T cells occurs in response to the peptide. T cell proliferation may be measured in several ways, e.g., by cellular uptake of tritiated thymidine. Stimulation indices for responses by T cells to peptides can be calculated as the maximum counts-per-minute (CPM) in response to a peptide divided by the control CPM. A stimulation index (S.I.) equal to or greater than two times the background level is considered "positive". Positive results are used to calculate the mean stimulation index for each peptide for the group of patients tested. Preferred peptides of this invention comprise at least one T cell epitope and have a mean T cell stimulation index of greater than or equal to 2.0. A peptide having a mean T cell stimulation index of greater than or equal to 2.0 in a significant number of ryegrass pollen sensitive patients tested (i.e., at least 10% of patients tested) is considered useful as a therapeutic agent. Preferred peptides have a mean T cell stimulation index of at least 2.5, more preferably at least 3.0, more preferably at least 3.5, more preferably at least 4.0, more preferably at least 5, and most preferably at least about 6. For example, peptides of the invention having a mean T cell stimulation index of at least 5, as shown in Fig. 3, include LPI-2 (SEQ ID NO: 5), LPI-3 (SEQ ID NO: 6), LPI-15 (SEQ ID NO: 21), LPI-16 (SEQ ID NO: 22), LPI-16.1 (SEQ ID NO: 23), LPI-17 (SEQ ID NO: 24), LPI-19 (SEQ ID NO: 26), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29) and LPI-23 (SEQ ID NO: 30). For example, peptides of the invention having a mean T cell stimulation index of at least 6, as shown in Fig. 3, include LPI-2 (SEQ ID NO: 5), LPI-15 (SEQ ID NO: 21), LPI-16 (SEQ ID NO: 22), LPI-16.1 (SEQ ID NO: 23), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), and LPI-23 (SEQ ID NO: 30).

In addition, preferred peptides have a positivity index (P.I.) of at least about 100, more preferably at least about 200 and most preferably at least about 300. The positivity index for a peptide is determined by multiplying the mean T cell stimulation index by the percent of individuals, in a population of individuals sensitive to ryegrass pollen (e.g., preferably at least 15 individuals, more preferably at least 30 individuals or more), who have a T cell stimulation index to such peptide of at least 2.0. Thus, the positivity index represents both the strength of a T cell response to a peptide (S.I.) and the frequency of a T cell response to a peptide in a population of individuals sensitive to ryegrass pollen. For example, as shown in Fig. 3, *Lol p I* peptide LPI-15 (SEQ ID NO: 21) has a mean S.I. of 12.2 and 11% of positive responses in the group of individuals tested resulting in a positivity index of 134.2. *Lol p I* peptides having a

positivity index of at least about 100 and a mean T cell stimulation index of at least about 4 include: LPI-2 (SEQ ID NO: 5), LPI-11 (SEQ ID NO: 15), LPI-13 (SEQ ID NO: 19), LPI-15 (SEQ ID NO: 21), LPI-16 (SEQ ID NO: 22), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), and LPI-23 (SEQ ID NO: 30).

In order to determine precise T cell epitopes by, for example, fine mapping techniques, a peptide having T cell stimulating activity and thus comprising at least one T cell epitope as determined by T cell biology techniques is modified by addition or deletion of amino acid residues at either the amino or carboxy terminus of the peptide and tested to determine a change in T cell reactivity to the modified peptide. If two or more peptides which share an area of overlap in the native protein sequence are found to have human T cell stimulating activity, as determined by T cell biology techniques, additional peptides can be produced comprising all or a portion of such peptides and these additional peptides can be tested by a similar procedure. Following this technique, peptides are selected and produced recombinantly or synthetically. Examples of fine map peptides are as follows: modified versions of peptide LPI-18 (SEQ ID NO: 25) (Fig. 2) include peptides: LPI-18.5 (SEQ ID NO: 39), LPI-18.6 (SEQ ID NO: 40), LPI-18.7 (SEQ ID NO: 41), LPI-18.8 (SEQ ID NO: 42) all as shown in Fig. 4; modified versions of peptide LPI-20 (SEQ ID NO: 27) (Fig. 2) include peptides: LPI-20.2 (SEQ ID NO: 43), LPI-20.3 (SEQ ID NO: 44), LPI-20.4 (SEQ ID NO: 45), LPI-20.5 (SEQ ID NO: 46), and LPI-20.6 (SEQ ID NO: 47) all as shown in Fig. 4; modified versions of peptide LPI-23 (SEQ ID NO: 30) (Fig. 2) include peptides: LPI-23.1 (SEQ ID NO: 48), LPI-23.2 (SEQ ID NO: 49) and LPI-23.4 (SEQ ID NO: 50) all as shown in Fig. 4.

Peptides are selected for diagnostic or therapeutic uses based on various factors, including the strength of the T cell response to the peptide (e.g., stimulation index), the frequency of the T cell response to the peptide in a population of individuals sensitive to ryegrass pollen, and the potential cross-reactivity of the peptide with other allergens from other species of grasses as discussed earlier. The physical and chemical properties of these selected peptides (e.g., solubility, stability) are examined to determine whether the peptides are suitable for use in therapeutic compositions or whether the peptides require modification as described herein. The ability of the selected peptides or selected modified peptides to stimulate human T cells (e.g., induce proliferation, lymphokine secretion) is determined.

The most preferred T cell epitope-containing peptides of the invention do not bind immunoglobulin E (IgE) of an allergic individual or bind IgE to a substantially lesser extent (e.g., at least 100 fold less and more preferably, at least 1000 fold less) than the protein allergen from which the peptide is derived. The major complications of standard immunotherapy are IgE-mediated responses such as anaphylaxis. Immunoglobulin E is a mediator of anaphylactic reactions which result from the binding and cross-linking of antigen to IgE on mast cells or basophils and the consequent release of mediators (e.g., histamine, serotonin, eosinophil chemotactic factors). Anaphylaxis in a substantial percentage of a population of individuals sensitive to *Lol p I* could be avoided by the use in immunotherapy of a peptide which do not bind IgE in a substantial percentage (e.g., at least about 75%) of a population of individuals sensitive to *Lol p I* allergen, or, if the peptides do bind IgE, such binding does not result in the release of mediators from mast cells or basophils. The risk of anaphylaxis could be reduced by the use in immunotherapy of a peptide or peptides which have reduced IgE binding. Moreover, peptides having minimal IgE stimulating activity are desirable for therapeutic effectiveness. Minimal IgE stimulating activity refers to IgE production that is less than the amount of IgE production stimulated by the native *Lol p I* protein allergen. Similarly, IL-4 production can be compared, with reduces IL-4 production indicating lessened IgE stimulating activity.

Preferred T cell epitope-containing peptides of the invention, when administered to a ryegrass pollen-sensitive individual or an individual sensitive to an allergen which is immunologically related to ryegrass pollen allergen (such as *Dac g I*, *Poa p I*, and *Phl p I*) in a therapeutic treatment regimen, are capable of modifying the allergic response of the individual to the allergen. Particularly, such preferred *Lol p I* peptides of the invention comprising at least one T cell epitope of *Lol p I* or at least two regions derived from *Lol p I*, each comprising at least one T cell epitope, when administered to an individual sensitive to ryegrass pollen are capable of modifying T cell response of the individual to the allergen, and they will thus be useful as therapeutics in addressing sensitivity to grasses.

A preferred isolated *Lol p I* peptide of the invention or portion thereof comprises at least one T cell epitope of *Lol p I* and accordingly, the peptide comprises at least approximately seven amino acid residues. For purposes of therapeutic effectiveness, preferred therapeutic compositions of the invention preferably comprise at least two T cell epitopes of *Lol p I*, and accordingly, the peptide comprises at least approximately eight amino acid residues and preferably at least fifteen amino acid

residues. Additionally, therapeutic compositions comprising preferred isolated peptides of the invention most preferably comprise a sufficient percentage of the T cell epitopes of the entire protein allergen so that a therapeutic regimen of administration of the composition to an individual sensitive to ryegrass pollen results in T cells of the individual being tolerized to the protein allergen. Synthetically produced peptides of the invention comprising up to approximately forty-five amino acid residues in length, and most preferably up to approximately thirty amino acid residues in length are particularly desirable, as increases in length may result in difficulty in peptide synthesis. Peptides of the invention may also be produced recombinantly as described above, and peptides exceeding 45 amino acids will be more easily produced recombinantly.

Peptides derived from the *Lol p I* protein allergen which exhibit T cell stimulatory properties and thus are believed to be useful therapeutics and/or intermediatea in developing tolerizing peptides comprise all or a portion of the following peptides: LPI-1 (SEQ ID NO: 3), LPI-1.1 (SEQ ID NO: 4), LPI-2 (SEQ ID NO: 5), LPI-3 (SEQ ID NO: 6), LPI-4 (SEQ ID NO: 7), LPI-4.1 (SEQ ID NO: 8), LPI-5 (SEQ ID NO: 9), LPI-6 (SEQ ID NO: 10), LPI-7 (SEQ ID NO: 11), LPI-8 (SEQ ID NO: 12), LPI-9 (SEQ ID NO: 13), LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-12 (SEQ ID NO: 17), LPI-13 (SEQ ID NO: 19), LPI-14 (SEQ ID NO: 20), LPI-15 (SEQ ID NO: 21), LPI-16 (SEQ ID NO: 22), LPI-16.1 (SEQ ID NO: 23), LPI-17 (SEQ ID NO: 24), LPI-18 (SEQ ID NO: 25), LPI-19 (SEQ ID NO: 26), LPI-20 (SEQ ID NO: 27), LPI-21 (SEQ ID NO: 28), LPI-22 (SEQ ID NO: 29), and LPI-23 (SEQ ID NO: 30) (Fig. 2) wherein the portion of the peptide preferably has a mean T cell stimulation index equivalent to, or greater than the mean T cell stimulation index of the corresponding peptide from which it is derived, as shown in Fig. 3. Even more preferably peptides derived from the *Lol p I* protein allergen comprise all or a portion of the following peptides: LPI-1.1 (SEQ ID NO: 4), LPI-2 (SEQ ID NO: 5), LPI-3 (SEQ ID NO: 6), LPI-4 (SEQ ID NO: 7), LPI-4.1 (SEQ ID NO: 8), LPI-8 (SEQ ID NO: 12), LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-13 (SEQ ID NO: 19), LPI-15 (SEQ ID NO: 21), LPI-16 (SEQ ID NO: 22), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), LPI-19 (SEQ ID NO: 26), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29) and LPI-23 (SEQ ID NO: 30), as shown in Fig. 2. Additionally, even more preferred peptides derived from the *Lol p I* protein comprise the following peptides: LPI-3 (SEQ ID NO: 6), LPI-4.1 (SEQ ID NO: 8), LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID

NO: 27), LPI-22 (SEQ ID NO: 29), and LPI-23 (SEQ ID NO: 30), all as shown in Fig. 2. Additional preferred peptides believed to T cell stimulating activity comprise the following peptides: LPI-16.2 (SEQ ID NO: 31), LPI-16.3 (SEQ ID NO: 32), LPI-16.4 (SEQ ID NO: 33), LPI-16.5 (SEQ ID NO: 34), LPI-16.6 (SEQ ID NO: 35), LPI-16.7 (SEQ ID NO: 36), LPI-16.9 (SEQ ID NO: 37), LPI-16.10 (SEQ ID NO: 38), LPI-18.5 (SEQ ID NO: 39), LPI-18.6 (SEQ ID NO: 40), LPI-18.7 (SEQ ID NO: 41), LPI-18.8 (SEQ ID NO: 42), LPI-20.2 (SEQ ID NO: 43), LPI-20.3 (SEQ ID NO: 44), LPI-20.4 (SEQ ID NO: 45), LPI-20.5 (SEQ ID NO: 46), LPI-20.6 (SEQ ID NO: 47), LPI-23.1 (SEQ ID NO: 48), LPI-23.2 (SEQ ID NO: 49), and LPI-23.4 (SEQ ID NO: 50).

One embodiment of the present invention features a peptide or portion thereof of *Lol p I* which comprises at least one T cell epitope of the protein allergen and has a formula X_n-Y-Z_m . According to the formula, Y is an amino acid sequence selected from the group consisting of LPI-1 (SEQ ID NO: 3), LPI-1.1 (SEQ ID NO: 4), LPI-2 (SEQ ID NO: 5), LPI-3 (SEQ ID NO: 6), LPI-4 (SEQ ID NO: 7), LPI-4.1 (SEQ ID NO: 8), LPI-5 (SEQ ID NO: 9), LPI-6 (SEQ ID NO: 10), LPI-7 (SEQ ID NO: 11), LPI-8 (SEQ ID NO: 12), LPI-9 (SEQ ID NO: 13), LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-12 (SEQ ID NO: 17), LPI-13 (SEQ ID NO: 19), LPI-14 (SEQ ID NO: 20), LPI-15 (SEQ ID NO: 21), LPI-16 (SEQ ID NO: 22), LPI-16.1 (SEQ ID NO: 23), LPI-17 (SEQ ID NO: 24), LPI-18 (SEQ ID NO: 25), LPI-19 (SEQ ID NO: 26), LPI-20 (SEQ ID NO: 27), LPI-21 (SEQ ID NO: 28), LPI-22 (SEQ ID NO: 29), LPI-23 (SEQ ID NO: 30), LPI-16.2 (SEQ ID NO: 31), LPI-16.3 (SEQ ID NO: 32), LPI-16.4 (SEQ ID NO: 33), LPI-16.5 (SEQ ID NO: 34), LPI-16.6 (SEQ ID NO: 35), LPI-16.7 (SEQ ID NO: 36), LPI-16.9 (SEQ ID NO: 37), LPI-16.10 (SEQ ID NO: 38), LPI-18.5 (SEQ ID NO: 39), LPI-18.6 (SEQ ID NO: 40), LPI-18.7 (SEQ ID NO: 41), LPI-18.8 (SEQ ID NO: 42), LPI-20.2 (SEQ ID NO: 43), LPI-20.3 (SEQ ID NO: 44), LPI-20.4 (SEQ ID NO: 45), LPI-20.5 (SEQ ID NO: 46), LPI-20.6 (SEQ ID NO: 47), LPI-23.1 (SEQ ID NO: 48), LPI-23.2 (SEQ ID NO: 49), and LPI-23.4 (SEQ ID NO: 50) and preferably selected from the group consisting of LPI-1.1 (SEQ ID NO: 4), LPI-2 (SEQ ID NO: 5), LPI-3 (SEQ ID NO: 6), LPI-4 (SEQ ID NO: 7), LPI-4.1 (SEQ ID NO: 8), LPI-8 (SEQ ID NO: 12), LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-13 (SEQ ID NO: 19), LPI-15 (SEQ ID NO: 21), LPI-16 (SEQ ID NO: 22), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), LPI-19 (SEQ ID NO: 26), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), LPI-23 (SEQ ID NO: 30), LPI-16.2 (SEQ ID NO: 31), LPI-16.3 (SEQ ID NO: 32), LPI-16.4 (SEQ ID NO: 33), LPI-16.5 (SEQ ID NO: 34), LPI-16.6 (SEQ ID NO: 35), LPI-16.7 (SEQ ID NO: 36), LPI-16.9 (SEQ ID NO: 37), LPI-16.10 (SEQ ID NO: 38), LPI-18.5 (SEQ ID NO: 39), LPI-18.6 (SEQ ID NO: 40), LPI-18.7 (SEQ ID NO: 41), LPI-18.8 (SEQ ID NO: 42), LPI-20.2 (SEQ ID NO: 43), LPI-20.3 (SEQ ID NO: 44), LPI-20.4 (SEQ ID NO: 45), LPI-20.5 (SEQ ID NO: 46), LPI-20.6 (SEQ ID NO: 47), LPI-23.1 (SEQ ID NO: 48), LPI-23.2 (SEQ ID NO: 49), and LPI-23.4 (SEQ ID NO: 50).

NO: 33), LPI-16.5 (SEQ ID NO: 34), LPI-16.6 (SEQ ID NO: 35), LPI-16.7 (SEQ ID NO: 36), LPI-16.9 (SEQ ID NO: 37), LPI-16.10 (SEQ ID NO: 38), LPI-18.5 (SEQ ID NO: 39), LPI-18.6 (SEQ ID NO: 40), LPI-18.7 (SEQ ID NO: 41), LPI-18.8 (SEQ ID NO: 42), LPI-20.2 (SEQ ID NO: 43), LPI-20.3 (SEQ ID NO: 44), LPI-20.4 (SEQ ID NO: 45), LPI-20.5 (SEQ ID NO: 46), LPI-20.6 (SEQ ID NO: 47), LPI-23.1 (SEQ ID NO: 48), LPI-23.2 (SEQ ID NO: 49), and LPI-23.4 (SEQ ID NO: 50) and more preferably selected from the group consisting of LPI-3 (SEQ ID NO: 6), LPI-4.1 (SEQ ID NO: 8), LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), LPI-23 (SEQ ID NO: 30), LPI-16.2 (SEQ ID NO: 31), LPI-16.3 (SEQ ID NO: 32), LPI-16.4 (SEQ ID NO: 33), LPI-16.5 (SEQ ID NO: 34), LPI-16.6 (SEQ ID NO: 35), LPI-16.7 (SEQ ID NO: 36), LPI-16.9 (SEQ ID NO: 37), LPI-16.10 (SEQ ID NO: 38), LPI-18.5 (SEQ ID NO: 39), LPI-18.6 (SEQ ID NO: 40), LPI-18.7 (SEQ ID NO: 41), LPI-18.8 (SEQ ID NO: 42), LPI-20.2 (SEQ ID NO: 43), LPI-20.3 (SEQ ID NO: 44), LPI-20.4 (SEQ ID NO: 45), LPI-20.5 (SEQ ID NO: 46), LPI-20.6 (SEQ ID NO: 47), LPI-23.1 (SEQ ID NO: 48), LPI-23.2 (SEQ ID NO: 49), and LPI-23.4 (SEQ ID NO: 50), and most preferably selected from the group consisting of LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO: 30), LPI-16.2 (SEQ ID NO: 31), LPI-16.3 (SEQ ID NO: 32), LPI-16.4 (SEQ ID NO: 33), LPI-16.5 (SEQ ID NO: 34), LPI-16.6 (SEQ ID NO: 35), LPI-16.7 (SEQ ID NO: 36), LPI-16.9 (SEQ ID NO: 37), LPI-16.10 (SEQ ID NO: 38), LPI-18.5 (SEQ ID NO: 39), LPI-18.6 (SEQ ID NO: 40), LPI-18.7 (SEQ ID NO: 41), LPI-18.8 (SEQ ID NO: 42), LPI-20.2 (SEQ ID NO: 43), LPI-20.3 (SEQ ID NO: 44), LPI-20.4 (SEQ ID NO: 45), LPI-20.5 (SEQ ID NO: 46), LPI-20.6 (SEQ ID NO: 47), LPI-23.1 (SEQ ID NO: 48), LPI-23.2 (SEQ ID NO: 49), and LPI-23.4 (SEQ ID NO: 50). In addition, X_n are amino acid residues contiguous to the amino terminus of Y in the amino acid sequence of the protein allergen and Z_m are amino acid residues contiguous to the carboxy terminus of Y in the amino acid sequence of the protein allergen. In the formula, n is 0-30 and m is 0-30. Preferably, the peptide or portion thereof has a mean T cell stimulation index equivalent to greater than the mean T cell stimulation index of Y as shown in Fig. 3. Preferably, amino acids comprising the amino terminus of X and the carboxy terminus of Z are selected from charged amino acids, i.e., arginine (R), lysine (K), histidine (H), glutamic acid (E) or aspartic acid (D); amino acids with reactive side chains, e.g., cysteine (C), asparagine (N) or glutamine (Q); or amino acids with sterically small side chains, e.g.,

alanine (A) or glycine (G). Preferably n and m are 0-5; most preferably n + m is less than 10.

Another embodiment of the present invention provides peptides comprising at least two regions, each region comprising at least one T cell epitope of *Lol p I* and accordingly each region comprises at least approximately seven amino acid residues. These peptides comprising at least two regions can comprise up to 100 or more amino acid residues but preferably comprise at least about 14, even more preferably at least about 20, and most preferably at least about 30 amino acid residues of the *Lol p I* allergen. If desired, the amino acid sequences of the regions can be produced and joined by a linker to increase sensitivity to processing by antigen-presenting cells. Such linker can be any non-epitope amino acid sequence or other appropriate linking or joining agent. To obtain preferred peptides comprising at least two regions, each comprising at least one T cell epitope, the regions are arranged in the same or a different configuration from a naturally-occurring configuration of the regions in the allergen. For example, the regions containing T cell epitope(s) can be arranged in a noncontiguous configuration and can preferably be derived from the same protein allergen. Noncontiguous is defined as an arrangement of regions containing T cell epitope(s) which is different than that of the native amino acid sequence of the protein allergen from which the regions are derived. Furthermore, the noncontiguous regions containing T cell epitopes can be arranged in a nonsequential order (e.g., in an order different from the order of the amino acids of the native protein allergen from which the region containing T cell epitope(s) are derived in which amino acids are arranged from an amino terminus to a carboxy terminus). A peptide of the invention can comprise at least 15%, at least 30%, at least 50% or up to 100% of the T cell epitopes of *Lol p I*.

The individual peptide regions can be produced and tested to determine which regions bind immunoglobulin E specific for *Lol p I* and which of such regions would cause the release of mediators (e.g., histamine) from mast cells or basophils. Those peptide regions found to bind immunoglobulin E and to cause the release of mediators from mast cells or basophils in greater than approximately 10-15% of the allergic sera tested are preferably not included in the peptide regions arranged to form preferred peptides of the invention.

Examples of preferred peptide regions which do not bind to IgE (data not shown) include: LPI-1 (SEQ ID NO: 3), LPI-1.1 (SEQ ID NO: 4), LPI-2 (SEQ ID NO: 5), LPI-3 (SEQ ID NO: 6), LPI-4 (SEQ ID NO: 7), LPI-4.1 (SEQ ID NO: 8),

LPI-5 (SEQ ID NO: 9), LPI-6 (SEQ ID NO: 10), LPI-7 (SEQ ID NO: 11), LPI-8 (SEQ ID NO: 12), LPI-9 (SEQ ID NO: 13), LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-12 (SEQ ID NO: 17), LPI-13 (SEQ ID NO: 19), LPI-14 (SEQ ID NO: 20), LPI-15 (SEQ ID NO: 21), LPI-16 (SEQ ID NO: 22), LPI-16.1 (SEQ ID NO: 23), LPI-17 (SEQ ID NO: 24), LPI-18 (SEQ ID NO: 25), LPI-19 (SEQ ID NO: 26), LPI-20 (SEQ ID NO: 27), LPI-21 (SEQ ID NO: 28), LPI-22 (SEQ ID NO: 29), LPI-23 (SEQ ID NO: 30), LPI-16.2 (SEQ ID NO: 31), LPI-16.3 (SEQ ID NO: 32), LPI-16.4 (SEQ ID NO: 33), LPI-16.5 (SEQ ID NO: 34), LPI-16.6 (SEQ ID NO: 35), LPI-16.7 (SEQ ID NO: 36), LPI-16.9 (SEQ ID NO: 37), LPI-16.10 (SEQ ID NO: 38), LPI-18.5 (SEQ ID NO: 39), LPI-18.6 (SEQ ID NO: 40), LPI-18.7 (SEQ ID NO: 41), LPI-18.8 (SEQ ID NO: 42), LPI-20.2 (SEQ ID NO: 43), LPI-20.3 (SEQ ID NO: 44), LPI-20.4 (SEQ ID NO: 45), LPI-20.5 (SEQ ID NO: 46), LPI-20.6 (SEQ ID NO: 47), LPI-23.1 (SEQ ID NO: 48), LPI-23.2 (SEQ ID NO: 49), and LPI-23.4 (SEQ ID NO: 50), the amino acid sequences of such regions being shown in Figs. 2 or 4, or portions of said regions comprising at least one T cell epitope.

Preferred peptides comprise various combinations of two or more of the above-discussed preferred regions, or a portion thereof. Preferred peptides comprising a combination of two or more regions (each region having an amino acid sequence as shown in Fig. 2 or Fig. 4), include the following:

- 20 LPI-3 (SEQ ID NO: 6), LPI-4.1 (SEQ ID NO: 8), LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), LPI-16 (SEQ ID NO: 22), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), and LPI-23 (SEQ ID NO: 30);
- 25 LPI-3 (SEQ ID NO: 6), LPI-4.1 (SEQ ID NO: 8), LPI-10 (SEQ ID NO: 14), and LPI-11 (SEQ ID NO: 15);
- LPI-3 (SEQ ID NO: 6), LPI-4.1 (SEQ ID NO: 8), LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), and LPI-16 (SEQ ID NO: 22);
- 30 LPI-3 (SEQ ID NO: 6), LPI-4.1 (SEQ ID NO: 8), LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), and LPI-16.1 (SEQ ID NO: 23);
- LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), and LPI-16.1 (SEQ ID NO: 23);

LPI-10 (SEQ ID NO:14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), and LPI-20 (SEQ ID NO: 27);

5 LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29) and LPI-23 (SEQ ID NO: 30);

LPI-15 (SEQ ID NO: 21), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), and LPI-20 (SEQ ID NO: 27);

10 LPI-15 (SEQ ID NO: 21), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), and LPI-23 (SEQ ID NO: 30);

LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), and LPI-23 (SEQ ID NO: 30);

LPI-18 (SEQ ID NO: 25) and LPI-20 (SEQ ID NO: 27);

15 LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27) and LPI-23 (SEQ ID NO: 30);

LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27) and LPI-16.1 (SEQ ID NO: 23);

20 LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO: 30) and LPI-16.1 (SEQ ID NO: 23);

LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO: 30), LPI-16.1 (SEQ ID NO: 23) and LPI-11 (SEQ ID NO: 15);

LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO: 30), LPI-16.1 (SEQ ID NO: 23) and LPI-4.1 (SEQ ID NO: 8);

25 LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO: 30), LPI-16.1 (SEQ ID NO: 23), LPI-4.1 (SEQ ID NO: 8) and LPI-22 (SEQ ID NO: 29);

LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO: 30), LPI-16.1 (SEQ ID NO: 23), LPI-11 (SEQ ID NO: 15) and LPI-4.1 (SEQ ID NO: 8);

30 LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO: 30), LPI-16.1 (SEQ ID NO: 23), LPI-11 (SEQ ID NO: 15), LPI-4.1 (SEQ ID NO: 8) and LPI-22 (SEQ ID NO: 29);

LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), and LPI-23 (SEQ ID NO: 30);

35

LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-16.1 (SEQ ID NO: 23), LPI-22 (SEQ ID NO: 29) and LPI-23 (SEQ ID NO: 30); and LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-16.1 (SEQ ID NO: 23) and LPI-22 (SEQ ID NO: 29).

5

Additional preferred peptides comprising various combinations of two or more of the above discussed preferred regions include:

- LPI-16.2 (SEQ ID NO: 31), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);
- 10 LPI-16.3 (SEQ ID NO: 32), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);
- LPI-16.4 (SEQ ID NO: 33), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);
- LPI-16.5 (SEQ ID NO: 34), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);
- 15 LPI-16.6 (SEQ ID NO: 35), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);
- LPI-16.7 (SEQ ID NO: 36), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);
- 20 LPI-16.9 (SEQ ID NO: 37), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30); and
- LPI-16.10 (SEQ ID NO: 38), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30).

25 In yet another aspect of the present invention, a composition is provided comprising at least two peptides (e.g., a physical mixture of at least two peptides), each comprising at least one T cell epitope of *Lol p I*. Such compositions can be in the form of a composition additionally with a pharmaceutically acceptable carrier of diluent for therapeutic uses, or with conventional non-pharmaceutical excipients for reagent use. When used therapeutically, an effective amount of one or more of such compositions can be administered simultaneously or sequentially to an individual sensitive to ryegrass pollen.

In another aspect of the invention, combinations of *Lol p I* peptides are provided which can be administered simultaneously or sequentially. Such combinations may comprise therapeutic compositions comprising only one peptide, or

35

more peptides if desired. Such compositions may be used simultaneously or sequentially in preferred combinations.

Preferred compositions and preferred combinations of *Lol p I* peptides which can be administered or otherwise used simultaneously or sequentially (comprising
 5 peptides having amino acid sequences shown in Fig. 2) include the following combinations:

- 10 LPI-3 (SEQ ID NO: 6), LPI-4.1 (SEQ ID NO: 8), LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), LPI-16 (SEQ ID NO: 22), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), and LPI-23 (SEQ ID NO: 30);
- LPI-3 (SEQ ID NO: 6), LPI-4.1 (SEQ ID NO: 8), LPI-10 (SEQ ID NO: 14), and LPI-11 (SEQ ID NO: 15);
- 15 LPI-3 (SEQ ID NO: 6), LPI-4.1 (SEQ ID NO: 8), LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), and LPI-16 (SEQ ID NO: 22);
- LPI-3 (SEQ ID NO: 6), LPI-4.1 (SEQ ID NO: 8), LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), and LPI-16.1 (SEQ ID NO: 23);
- 20 LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), and LPI-16.1 (SEQ ID NO: 23);
- LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), and LPI-20 (SEQ ID NO: 27);
- 25 LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29) and LPI-23 (SEQ ID NO: 30);
- LPI-15 (SEQ ID NO: 21), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), and LPI-20 (SEQ ID NO: 27);
- 30 LPI-15 (SEQ ID NO: 21), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), and LPI-23 (SEQ ID NO: 30);
- LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), and LPI-23 (SEQ ID NO: 30);
- LPI-18 (SEQ ID NO: 25) and LPI-20 (SEQ ID NO: 27);

- LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27) and LPI-23 (SEQ ID NO: 30);
- LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27) and LPI-16.1 (SEQ ID NO: 23);
- 5 LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO: 30) and LPI-16.1 (SEQ ID NO: 23);
- LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO: 30), LPI-16.1 (SEQ ID NO: 23) and LPI-11 (SEQ ID NO: 15);
- 10 LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO: 30), LPI-16.1 (SEQ ID NO: 23) and LPI-4.1 (SEQ ID NO: 8);
- LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO: 30), LPI-16.1 (SEQ ID NO: 23), LPI-4.1 (SEQ ID NO: 8) and LPI-22 (SEQ ID NO: 29);
- 15 LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO: 30), LPI-16.1 (SEQ ID NO: 23), LPI-11 (SEQ ID NO: 15) and LPI-4.1 (SEQ ID NO: 8);
- LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO: 30), LPI-16.1 (SEQ ID NO: 23), LPI-11 (SEQ ID NO: 15), LPI-4.1 (SEQ ID NO: 8) and LPI-22 (SEQ ID NO: 29);
- 20 LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), and LPI-23 (SEQ ID NO: 30);
- LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-16.1 (SEQ ID NO: 23), LPI-22 (SEQ ID NO: 29) and LPI-23 (SEQ ID NO: 30); and
- 25 LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-16.1 (SEQ ID NO: 23) and LPI-22 (SEQ ID NO: 29).

Additional preferred compositions and preferred combinations of *Lol p I* peptides which can be administered or used simultaneously or sequentially (comprising peptides having amino acid sequences shown in Figs. 2 or 4) include the following

30 combinations:

- LPI-16.2 (SEQ ID NO: 31), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);
- LPI-16.3 (SEQ ID NO: 32), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);

LPI-16.4 (SEQ ID NO: 33), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);
 LPI-16.5 (SEQ ID NO: 34), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);
 5 LPI-16.6 (SEQ ID NO: 35), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);
 LPI-16.7 (SEQ ID NO: 36), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);
 LPI-16.9 (SEQ ID NO: 37), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30); and
 10 LPI-16.10 (SEQ ID NO: 38), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30).

In each of the above preferred compositions, peptides LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 23), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30) may be substituted as follows: peptide LPI-16.1 (SEQ ID NO: 23) (Fig. 2) may be substituted with LPI-16.2 (SEQ ID NO: 31), LPI-16.3 (SEQ ID NO: 32), LPI-16.4 (SEQ ID NO: 33), LPI-16.5 (SEQ ID NO: 34), LPI-16.6 (SEQ ID NO: 35), LPI-16.7 (SEQ ID NO: 36), LPI-16.9 (SEQ ID NO: 37), and LPI-16.10 (SEQ ID NO: 38), all as shown in Fig. 4; peptide LPI-18 (SEQ ID NO: 25) (Fig. 2) may be substituted with peptides LPI-18.5 (SEQ ID NO: 39), LPI-18.6 (SEQ ID NO: 40), LPI-18.7 (SEQ ID NO: 41), LPI-18.8 (SEQ ID NO: 42) all as shown in Fig. 4; peptide LPI-20 (SEQ ID NO: 27) may be substituted with peptides LPI-20.2 (SEQ ID NO: 43), LPI-20.3 (SEQ ID NO: 44), LPI-20.4 (SEQ ID NO: 45), LPI-20.5 (SEQ ID NO: 46), and LPI-20.6 (SEQ ID NO: 47) all as shown in Fig. 4; peptide LPI-23 (SEQ ID NO: 30) may be substituted with peptides LPI-23.1 (SEQ ID NO: 48), LPI-23.2 (SEQ ID NO: 49) and LPI-23.4 (SEQ ID NO: 50), all as shown in Fig. 4.

The present invention is further illustrated by the following non-limiting Figures and Examples.

30

EXAMPLES

Example 1 - Isolation and Cloning of Nucleic Acid Sequence Coding for *Lol p I*

Total mRNA was extracted from mature ryegrass pollen by the phenol method of Herrin and Michaels, *supra*. Double-stranded cDNA was synthesized from 1µg of

35

total mRNA using a commercially available kit (cDNA SYNTHESSES SYSTEM PLUS KIT, BRL, Gaithersburg, MD). After a phenol extraction and ethanol precipitation, the cDNA was blunted with T4 DNA polymerase (Promega, Madison, WI) and ligated to ethanol-precipitated, self-annealed AT and AL oligonucleotides for use in a
5 modified Anchored PCR reaction, according to the method in Rafnar *et al.* (1991), *J. Biol. Chem.*, **266**: 1229-1236; Frohman *et al.* (1990), *Proc. Natl. Acad. Sci. USA*, **85**:8998-9002; and Roux *et al.* (1990), *BioTech.*, **8**: 48-57. Oligonucleotide AT has the sequence 5'-GGGTCTAGAGGTACCGTCCGATCGATCATT-3' (SEQ ID NO: 71) (Rafnar *et al. supra*). Oligonucleotide AL has the sequence
10 AATGATCGATGCT (SEQ ID NO: 72) (Rafnar *et al. supra*).

Polymerase chain reactions (PCR) were carried out using a commercially available kit (GeneAmp® DNA Amplification kit, Perkin Elmer Cetus, Norwalk, CT) whereby 10 µl 10x buffer containing dNTPs were mixed with 1 µg each of primer AP, which has the sequence 5'-GGGTCTAGAGGTACCGTCCG-3' (SEQ ID NO: 73)
15 (Rafnar *et al. supra*) and LpA-5, which has the sequence 5'-CCCTGCAGATTATTTGAGATCTTGAG-3' (SEQ ID NO: 74), cDNA (3-5 µl of a 20 µl linkered cDNA reaction mix), 0.5 µl Amplitaq DNA polymerase, and distilled water to 100 µl.

Nucleotides 1 through 8 (5'-CCCTGCAG) of LpA-5 correspond to a Pst I site
20 added for cloning purposes; the remaining nucleotides correspond to the non-coding strand sequence complementary to nucleotides 483 through 500 as shown in Fig. 6.

The samples were amplified with a programmable thermal controller (MJ Research, Inc., Cambridge, MA). The first 5 rounds of amplification consisted of denaturation at 94°C for 1 minute, annealing of primer to the template at 45°C for 1.5
25 minutes, and chain elongation at 70°C for 2 minutes. The final 20 rounds of amplification consisted of denaturation as above, annealing at 55°C for 1.5 minutes, and elongation as above. Five percent (5 µl) of this initial amplification was then used in a secondary amplification whereby 10 µl 10x buffer containing dNTPs was mixed with 1 µg each of primer AP and primer LpA-3, which has the sequence 5'-
30 CCCTGCAGTCATGCTCACTTGGCCGAGTA-3' (SEQ ID NO: 75), 0.5 µl Amplitaq DNA polymerase, and distilled water to 100 µl. The secondary PCR reaction was performed as described herein. Nucleotides 1 through 8 (5'-CCCTGCAG-3') of LpA-3 correspond to a Pst I site added for cloning purposes; nucleotides 9 through 12 (5'-TCA-3') correspond to the complementary sequence for a new stop codon, and the
35 remaining nucleotides correspond to the non-coding strand sequence complementary

to nucleotides 793 through 810 of the full length clone of *Lol p I* as shown in Fig. 1, including translated sequence of *Lol p I* (Fig. 1), the native stop codon and 3' untranslated sequence.

Amplified DNA was recovered by sequential chloroform, phenol, and
 5 chloroform extractions, followed by precipitation at -20°C with 0.5 volumes of 7.5 ammonium acetate and 1.5 volumes of isopropanol. After precipitation and washing with 70% ethanol, the DNA was simultaneously digested with *Xba I* and *Pst I* in a 15 µl reaction and electrophoresed through a preparative 3% GTG NuSieve low melt gel (FMC, Rockport, ME). The appropriate sized DNA band was visualized by EtBr
 10 staining, excised, and ligated into appropriately digested M13mp18 for sequencing by the dideoxy chain termination method (Sanger *et al.* (1977), *Proc. Natl Acad Sci USA*, 74: 5463-5476) using a commercially available sequencing kit (Sequenase kit, U.S. Biochemicals, Cleveland, OH).

Both strands were sequenced using M13 forward and reverse primers (N.E.
 15 BioLabs, Beverly, MA) and internal sequencing primers LpA-13, LpA-12, LpA-9, LpA-2, LpA-7, LpA-10, and LpA-IA. LpA-13 has the sequence 5'-GAGTACGGCGACAAGTGGC-3' (SEQ ID NO: 76), which corresponds to nucleotides 121 through 139 as shown in Fig. 1. LpA-12 has the sequence 5'-TTCGAGATCAAGTGCACC-3' (SEQ ID NO: 77), which corresponds to nucleotides
 20 310 through 318 as shown in Fig. 1. LpA-9 has the sequence 5'-GTGACAGCCTCGCCGG-3' (SEQ ID NO: 78), which corresponds to the non-coding strand sequence complementary to nucleotides 335 through 350 as shown in Fig. 1. LpA-2 has the sequence 5'-GGGAATTCCATGGCGAAGAAGGGC-3' (SEQ ID NO: 79). Nucleotides 1 through 7 (5-GGGATT-3') of LpA 2 correspond to part
 25 of an Eco-RI restriction site added for cloning purposes; the remaining sequence of LpA-2 corresponds to nucleotides 425 through 441 of Fig. 1. LpA-7 has the sequence 5'-GTGCCGTCCGGGTACT-3' (SEQ ID NO: 80), and corresponds to non-coding strand sequence complementary to nucleotides 503 through 518 of Fig. 1. LpA-10 has the sequence 5'-CCGTCGACGTACTTCA-3' (SEQ ID NO: 81), which corresponds to
 30 non-coding strand sequence complementary to nucleotides 575 through 590 of Fig. 1. LpA-IA has the sequence 5'-GGAGTCGTGGGGAGCAGTC-3' (SEQ ID NO: 82), which corresponds to nucleotides 654 through 672 of Fig. 1.

Multiple clones from several independent PCR reactions were sequenced. The
 nucleotide (SEQ ID NO: 1) and deduced amino acid sequences (SEQ ID NO: 2) of a
 35 representative clone of *Lol p I*, clone 26.j are shown in Fig. 1. As shown in Fig.1, the

nucleic acid sequence coding for *Lol p I* has an open reading frame beginning with an ATG initiation codon at nucleotides 16-18 ending with a TGA stop codon at nucleotides 805-807. The translated protein has a deduced amino acid sequence of 263 amino acids with a predicted molecular weight of 28.4 kD and a pI of 5.55. The
5 initiating methionine is numbered amino acid -23, with amino acid numbered +1 corresponding to the NH₂-terminus of the mature protein, as defined by amino acid sequencing (Cottam *et al.* (1986), *Biochem. J.*, **234**: 305-310). Amino acids -23 through -1 (Fig.1), correspond to a leader sequence that is cleaved to yield the mature protein; the mature protein is therefore composed of 240 amino acids and has a
10 predicted molecular weight of 26.1 kD and a pI of 5.38. There is a single potential N-linked glycosylation site at amino acid 9.

Amino acids 1 through 30 of clone 26.j (Fig. 1) correspond exactly to the published sequence of the NH₂ terminus of *Lol p I* (Cottam *et al.*, *supra*). Amino acids 213 through 240 of clone 26.j (Fig. 1) correspond exactly to the published internal
15 amino acid sequence of *Lol p I* (Esch and Klapper (1989), *Mol. Immunol.*, **26**: 557-561).

Example 2 - Identification of Polymorphisms in *Lol p I*

A number of polymorphisms in the nucleotide sequence coding for *Lol p I* were
20 discovered during the amplification and sequencing of different *Lol p I* clones. Some of the polymorphisms cause an amino acid change relative to that of clone 26.j, while others are silent polymorphisms that do not cause an amino acid change. The polymorphisms found in the sequence coding for *Lol p I* are summarized in Table 1. The nucleotide base numbers are those of the sequence of clone 26.j shown in Fig 1.

Table 1**Polymorphisms Detected in *Lol p I***

5	Nucleotide Polymorphism	Amino Acid Polymorphism
1	GGC ₂₁₅ →GGA/GGT	None
2	G ₂₃₄ AC ₂₃₆ →GAT	D ₄₅ →N
3	GTT ₂₃₉ →GTC	None
4	CGT ₃₅₁ →CGC	None
5	GGC ₃₅₆ →GGT	None
6	AAC ₃₈₉ →AAT	None
7	CCC ₃₉₆ →CCT	None
8	CAT ₄₁₃ →CAC	None
9	GCC ₄₃₄ →GCA	None
10	GAC ₅₃₀ →GAT	None
11	GG ₅₃₂ C→GAC	G ₁₄₄ →D
12	CCG ₅₄₂ →CCA	None
13	ACA ₅₄₅ →ACG	None
14	GC ₅₆₂ T→GGT	A ₁₅₄ →G
15	CTC ₅₈₁ →CTG	None
16	GCG ₆₂₆ →GCC	None
17	ATC ₇₈₂ →ATT	None
18	CCT ₇₈₅ →CCC	None

All confirmed nucleotide polymorphisms (polymorphisms observed in the sequence analysis of clones from two independent PCR reactions) are shown relative to the sequence of clone 26.j (Fig.1) (SEQ ID NO: 1). The polymorphic residues in their respective codon triplets are numbered. Productive amino acid changes are also shown; most nucleotide polymorphisms are silent and do not result in an amino acid change. Twenty-eight potential polymorphisms have only been observed in clones from single PCR reactions. Seventeen of these 28 potential polymorphisms are silent mutations and do not result in an amino acid polymorphism; the remaining 11 potential polymorphic sites would result in the following amino acid changes, specifically: T₁₁

→M, A49→V, R67→S, K79→R, V90→I, Q133→R, I162→T, V173→E, I187→T, V223→F and K232→R. The potential polymorphism at amino acid 223 (V223→F) has been previously reported. (Perez *et al.*, *supra*)

5 **Example 3 - Human IgE reactivity to Purified Recombinant and Native *Lol p I***

Cloned DNA encoding *Lol p I* and *Lol p IX* was expressed in *E. coli* and purified on a Ni-chelating affinity column. Monoclonal antibodies were also used to affinity purify and distinguish isoforms of these and native grass proteins. The recombinant *Lol p I* was compared to biochemically purified native *Lol p I* and *Lol p*
10 *IX* in mAb and human IgE reactivity studies (data not shown). The reactivity of human IgE to the recombinant and native forms was equivalent when measured by direct binding ELISA. In competition assays, the native *Lol p I* and *Lol p IX* proteins could completely inhibit IgE binding to *Lol p* soluble pollen extract (SPE), whereas the recombinant form of *Lol p I* and *Lol p IX* could only partially inhibit IgE binding to the
15 extract. However, the recombinant *Lol p I* and *Lol p IX* was still active in these competition assays. These assays were then extended to western blot inhibition studies; both methods confirm the previous finding that Group I and Group IX constitute one of the major allergenic proteins of *Lolium perenne* grass pollen. Furthermore, the *Lol p I* and *Lol p IX* native and recombinant allergens showed inhibition of grass allergic
20 patient IgE binding to soluble pollen extracts of other grass species (*Dac g*, *Phl p* and *Poa p*). The degree to which *Lol p I* and *Lol p IX* proteins successfully compete for IgE binding to these other grasses implies a hierarchy of homology between the species. These studies confirm and extend the findings of shared IgE epitopes between temperate grass allergens.

25 The procedures used for the foregoing examples were as follows:

Extraction and Depigmentation of Allergens

Defatted *Lol p I* pollen was extracted twice, overnight at 4°C in 50mM phosphate buffer, 15mM NaCl, pH 7.2 and protease inhibitors (PMSF, Luepeptin, SPTI and pepstatin). The extract was then depigmented by batch absorption with DE-
52 (Whatman) in 50mM phosphate buffer, 0.3M NaCl, pH 7.2.

Biochemical Purification of *Lol p I* Allergen

Depigmented *Lol p I* extract was dialyzed into H₂O, pH 8.0 by addition of NH₄OH. This material was loaded onto a DE-52 column and eluted stepwise with
1mM, 4.5mM and 7.5mM NaH₂PO₄. The majority of the Group I allergens was eluted with 4.5mM NaH₂PO₄. A further separation of Group I was accomplished by running this DE-52 enriched fraction over A (26/60) superdex 75 column (Pharmacia).

Immunoaffinity Purification of *Lol p IX* Allergen

1B9 ascites was precipitated by 50% (NH₄)₂SO₄, followed by purification over Q-sepharose (Pharmacia). Purified 1B9, an anti-*Lol p IX* antibody, was then coupled to Affigel-10 (Biorad), according to the manufacturer's instructions. Either depigmented pollen extract or DE-52 enriched material was circulated over the 1B9 affigen column overnight at 4°C. The column was washed with PBS, PBS + 0.5M
MaCl and then eluted with 0.1M Glycine, pH2.7. Eluted *Lol p IX* fractions were neutralized with 1M tris-base, pH 11.

Expression and Purification of Recombinant *Lol p I*

Lol p I cDNA's encoding from the first amino acid of the mature protein to the stop codon were ligated into pET11dΔHR containing a leader which encoded 6 histidines. The HIS₆ was used for purification over a nickel-NTA agarose column (Qiagen). r*Lol p I* was expressed in *E. coli*.

SDS-PAGE, Electrophoresis and Immunoblotting

Electrophoresis was performed using 12.5% polyacrylamide gels. The samples were run under reducing conditions (4 hours at 40mA constant current). After electrophoresis the protein was transferred to nitrocellulose membrane (1.5 hours at 1.5A). The blots were stained with 1% India ink, and then blocked with 1% defatted milk, 1% FCS in Tween solution (2mM Tris-HCl pH 7.5, 0.71M NaCl, and 0.05% Tween 20) for 1 hour. The human plasma samples were pre-absorbed with blank

nitrocellulose for 1.5 hours prior to incubation. Blot sections were incubated with 1st antibodies diluted in 1% milk/Tween solution overnight at room temperature (RT). The blot sections were washed three times and incubated in the appropriate biotinylated 2nd AB (1:2500) for 2 hours at RT. The blot sections were washed three times and finally incubated with ^{125}I -streptavidin 1 hour at RT. The sections were washed extensively to remove unbound label and exposed to film. Autoradiography was carried out at -80°C .

Direct, Competition and Depletion ELISA

Microtiter plates were coated with 2.5-10.0 $\mu\text{g/mL}$ of coating antigen (grass soluble pollen extract (SPE), *Lol p I*, *Lol p IX*, *Lol p IX*, recombinant *Lol p I*, and/or recombinant *Lol p IX*) in PBS at 100 μL /well and incubated overnight at 4°C . The plates were washed three times between each step with PBS-T (Phosphate buffered saline + 0.05% Tween 20). The unbound antigen was removed and the plate blocked with 300 μL /well of 1MG/ML PVP in 0.5% gelatin/PBS for one hour at room temperature (RT). All subsequent reagents were added at 100 μL /well for direct ELISA, serially diluted human plasma was added to duplicate wells and incubated overnight at 4°C . This was followed by biotinylated goat anti-human IgE (1:1,000) for 1 hour at RT, then streptavidin-HRPO (1:10,000) for 1 hour at RT. TMB substrate and H_2O_2 were freshly mixed and added; the color was allowed to develop for 2-5 minutes. The reaction was stopped by the addition of 1M phosphoric acid. The plates were read on a dynatech plate reader at 450NM and the absorbances of duplicate wells were averaged.

For the competition ELISA, the human plasma samples were mixed with an equal volume of serially diluted antigen or with PBS-T (as a control). These samples were incubated overnight at 4°C before addition to the microtiter plate and performing the remaining steps of the ELISA as stated above.

For the depletion ELISA, the human plasma was pre-incubated on antigen or PBS coated wells, collected and re-incubated on freshly coated wells. The ELISA was then performed as outlined above.

EXAMPLE 4 - Human T Cell Studies with *Lol p I*

Synthesis of Overlapping Peptides

Ryegrass *Lol p I* overlapping peptides were synthesized using standard
5 Fmoc/tBoc synthetic chemistry and purified by Reverse Phase HPLC. Fig. 2 shows
Lol p I peptides used in these studies (SEQ ID NO: 3-30). The peptide names are
consistent throughout.

IgE Binding Studies with overlapping peptides

10 None of the peptides shown in Fig. 2 bound a detectable amount of IgE from
pooled human plasma when analyzed in a solid phase ELISA assay (data not shown).
The procedure for the ELISA assay with the overlapping peptides was substantially the
same as that described in Example 3.

T Cell Responses to Ryegrass Antigen Peptides

Peripheral blood mononuclear cells (PBMC) were purified by lymphocyte
separation medium (LSM) centrifugation of 60 ml of heparinized blood from grass-
allergic patients who exhibited clinical symptoms of seasonal rhinitis and were MAST
and/or skin test positive for grass. Long-term T cell lines were established by
20 stimulation of 2×10^6 PBL/ml in bulk cultures of complete medium IRPMI-1640, 2 mM
L-glutamine, 100 U/ml penicillin/streptomycin, 5×10^{-5} M 2-mercaptoethanol, and 10
mM HEPES, supplemented with 5% heat-inactivated human AB serum) with 25
mg/ml of purified native *Lol p I* (95% pure with a single band on protein gel) for 6
days at 37°C in a humidified 5% CO₂ incubator to select for *Lol p I* reactive T Cells.
25 This amount of priming antigen was determined to be optimal for the activation of T
cells from most grass-allergic patients. Viable cells were purified by LSM
centrifugation and cultured in complete medium, supplemented with 5 units
recombinant human IL-2/ml and 5 units recombinant human IL-4/ml for up to 3 weeks
until the cells no longer responded to lymphokines and were considered "rested." The
30 ability of the T cells to proliferate to selected peptides, recombinant *Lol p I* (r*Lol p I*),
purified native *Lol p I*, recombinant *Lol p IX* (r*Lol p IX*), or *Der p I* (r*Der p I*) was
then assessed. For assay, 2×10^4 rested cells were restimulated in the presence of
 2×10^4 autologous Epstein-Barr virus (EBV)-transformed B cells (prepared as
described below) with 2-50 mg/ml of r*Lol p I*, purified native *Lol p I*, r*Der p I*, or r*Lol*
35 *p IX*, in a volume of 200 µl complete medium in duplicate wells in 96-well round-

bottom plates for three days. Each well then received 1 mCi tritiated thymidine for 16-20 hours. The counts incorporated were collected onto glass fiber filter mats and processed for liquid scintillation counting. The varying antigen dose in assays with *rLol p I*, purified native *Lol p I*, and recombinant *Lol p IX* and several antigenic peptides (i.e., peptides that induce an immune response, or, specifically, a positive T cell response in these assays) synthesized as described above were determined. The titrations were used to optimize the dose of peptides in T cell assays. The maximum response in a titration of each peptide is expressed as the stimulation index (S.I.). The S.I. is the counts per minute (CPM) incorporated by cells in response to peptide, divided by the CPM incorporated by cells in medium only. An S.I. value equal to or greater than 2 times the background level is considered "positive" and indicates that the peptide contains a T cell epitope. The positive results were used in calculating mean stimulation indices for each peptide for the group of patients tested. The results (not shown) demonstrate that one patient responds well to *rLol p I* and purified native *Lol p I*, as well as to *Lol p I* peptides but not to recombinant *Der p I*. This indicated that *Lol p I* T cell epitopes are recognized by T cells from this particular allergic patient and that *rLol p I* contains such T cell epitopes. T cells from the majority of patients also reacted to *rLol p IX*, suggesting a presence of *Lol p IX* antigen in the purified native *Lol p I* prep that was used to prime T cells.

The above procedure was followed with a number of other patients. Individual patient results were used in calculating the mean S.I. for each peptide if the patient responded to the *Lol p I* protein at an S.I. of 2.0 or greater and the patient responded to at least one peptide derived from *Lol p I* at an S.I. of 2.0 or greater. A summary of positive experiments from 35 patients is shown in Fig. 3. All 35 T cell lines responded to purified native *Lol p I* and *rLol p I*. The numbers enclosed in the parentheses denote percentage of patients responding to that particular peptide. The bar represents the positivity index for each peptide (% of patients responding multiplied by mean S.I.).

Preparation of EBV-transformed B Cells for Use as Antigen-presenting Cells

Autologous EBV-transformed cell lines were derived by incubating 5×10^6 PBL with 1 ml of B-59/8 Marmoset cell line (ATCC CRL1612, American Type Culture Collection, Rockville, MD) conditioned medium in the presence of 1 mg/ml phorbol 12-myristate 13-acetate (PMA) at 37°C for 60 minutes in 12x75 mm polypropylene round-bottom Falcon snap cap tubes (Becton Dickinson Labware, Lincoln Park, NJ).

These cells were then diluted to 1.25×10^6 cells/ml in the RPMI-1640 medium that was supplemented with 10% head-inactivated fetal bovine serum in place of the 5% human AB serum and cultured in 200 ml aliquots in flat-bottom culture plates until visible colonies were detected. They were then transferred to larger wells until the cell lines were established.

Those skilled in the art will appreciate that the invention described is susceptible to variations and modification other than those specifically described. It is understood that the invention includes all such variations and modifications. The invention also includes all steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all combinations of any two or more of said steps or features.

Example 5 - Cloning and Expression of *Dac g I*, *Poa p I* and *Phl p I*

15 A. Cloning of *Dac g 1*.

RNA was obtained from the pollen of *Dactylis glomerata* using a standard acid phenol extraction procedure (Sambrook *et al.* (1989), *Molecular Cloning: A laboratory manual. 2nd Edition.*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY). This and other pollens described below were purchased from Greer Laboratories (Lenoir, NC). Single and double stranded cDNA was prepared from total *D. glomerata* RNA using the BRL cDNA Synthesis System (Gaithersburg, MD), blunted using standard procedures (Sambrook *et al.* (1989) *supra*), and ligated to self-annealed oligonucleotides AT (5'-GGGTCTAGAGGTACCGTCCGATCGATCATT-3') (SEQ ID NO: 71) and AL (5'-AATGATCGATGCT-3') (SEQ ID NO: 72) (Rafnar *et al.* (1991), *J. Biol. Chem.*, **266**:1229-1236).

The amino portion of the gene encoding *Dac g 1*, including 5' untranslated sequence, nucleotide sequence encoding the predicted leader sequence and nucleotide sequence encoding the first portion of the mature protein, was cloned using the polymerase chain reaction (PCR). Oligonucleotide primers AP-2 (5'-GGGTCTAGAGGTACCGTCC-3') (SEQ ID NO: 83) and LpA-7 (5'-GTGCCGTCCGGGTACT-3') (SEQ ID NO: 80) were used in a primary amplification. Oligonucleotide primers AP-2 and LpA-9 (5'-GTGACAGCCTCGCCGG-3') (SEQ ID NO: 78) were used in a secondary amplification using 10% of the primary amplification as template cDNA. PCRs were carried out using the GeneAmp DNA Amplification kit (Perkin Elmer, Norwalk, CT)

using a programmable thermal controller from MJ Research, Inc. (Cambridge, MA). Samples were amplified for 24 cycles by heating to 94°C for 1 min, 54°C for 1.5 min and 70°C for 1 min.

The resulting PCR product was blunted with T4 DNA polymerase (Sambrook *et al.* (1989) *supra*) and digested with the restriction endonuclease *Xba*I. Unless
5 otherwise stated, all endonucleases and polymerases were obtained from New England BioLabs (Beverly, MA). A band of approximately 400 base pairs was isolated from a low melting temperature agarose gel (FMC, Rockland, ME) and ligated into appropriately digested pUC19. The clones 22.2 and 22.5 were subsequently identified
10 by dideoxysequencing (Sanger *et al.* (1977), *Proc. Natl. Acad. Sci. USA*, **74**:5460-5463) to contain nucleotide sequence of the gene encoding *Dac g 1*.

A 600 base pair cDNA containing internal nucleotide sequence of the gene encoding *Dac g 1* was amplified using the primers DGI-3 (5'-
TTGGATCCTACGGCAAGCCGACCGGC-3') (SEQ ID NO: 84) and LpA-10 (5'-
15 CCGTCGACGTACTTCA-3') (SEQ ID NO: 81). A 300 base pair cDNA containing internal *Dac g 1* sequence was amplified using the primers DGI-4 (5'-
TTGGATCCATCCCGAAGGTGCCCCCGGG-3' (SEQ ID NO: 85), wherein G at position 14 can also be A) and LpA-9 (5'-GTGACAGCCTCGCCGG-3') (SEQ ID NO: 78). The cDNAs were amplified for 34 cycles by heating to 94°C for 45 sec,
20 60°C for 45 sec and 72°C for 1 min. These PCR products were blunted with T4 DNA polymerase as above, digested with *Bam*HI and ligated into appropriately digested pUC19. Clones 86.1 (600 base pairs) and 88.6 (300 base pairs) were sequenced and found to contain sequence of the gene encoding *Dac g 1*.

The carboxy portion of the gene encoding *Dac g 1*, including the 3'
25 untranslated region, was cloned using oligonucleotide primers AP (5'-GGGTCTAGAGGTACCGTCCG-3') (SEQ ID NO: 73) and DGI-8 (5'-AGGTGACCTTCCACGTCG-3') (SEQ ID NO: 86) in a primary PCR and oligonucleotide primers AP and DGI-9 (5'-
TTGGATCCTGGCGCTGCTGGTGAAGTA-3') (SEQ ID NO: 87) in a secondary
30 PCR. Material was amplified for 25 cycles of heating to 94°C for 1 min, 60°C for 40 sec and 74°C for 1 min. The 700 base pair PCR product was digested with *Bam*HI and *Asp*718 (Boehringer Mannheim, Indianapolis, IN), isolated and digested into appropriately digested pUC19 as described above. The clones 119.2, 119.4, 119.6, 119.9 and 119.12 were isolated, sequenced and found to contain sequence of the gene
35 encoding *Dac g 1*.

cDNA clones encoding the mature *Dac g 1* protein were obtained by PCR with the oligonucleotide primers DGI-7Eco (5'-

TTGAATTCATCCCGAAGGTGCCCCCG-3' (SEQ ID NO: 88), wherein G at position 14 can also be A) and PhA-1.2 (5'-

- 5 TTGGTACCTCACTTGGACTCGTAGCT-3') (SEQ ID NO: 89). The cDNAs were amplified for 24 cycles of heating to 94°C for 1 min, 54°C for 1.5 min and 70°C for 1 min. The amplified cDNA was digested with *EcoRI* and *Asp718*, isolated, and ligated into the appropriately digested pUC19. The cDNA clones 106.5, 106.6, 106.9 and 106.12 were identified as containing *Dac g 1* sequence by dideoxysequencing. The
 10 nucleotide (SEQ ID NO: 51) and deduced amino acid (SEQ ID NO: 52) sequences of clone 106.5 are shown in Fig. 5. Nucleotides 509-515 (encoding amino acids 171 and 172) are from the sequence of clone 106.12. The sequence of clone 106.5 was not resolved in this region.

- The insert from clone 106.5 was isolated and ligated into appropriately
 15 digested expression vector pET-11d (Novagen, Madison, WI: Jameel et al. (1990), *J. Virol.*, **64**:3963-3966). The pET-11d vector had been modified to contain a sequence encoding 6 histidines (His 6) immediately 3' of the ATG initiation codon followed by a unique *EcoRI* endonuclease restriction site. A second *EcoRI* endonuclease restriction site in the vector, along with neighboring *ClaI* and *HindIII* endonuclease restriction
 20 sites, had previously been removed by digestion with *EcoRI* and *HindIII*, blunted and religated.

- A recombinant clone was used to transform *Escherichia coli* strain BL21-DE3. A culture was grown to A₆₀₀ of 1.0, IPTG was added to 1 mM final concentration and grown for an additional 2 hours. Bacteria was recovered by
 25 centrifugation (7,930 G, 10 min) and lysed in 90 ml of 6 M Guanidine-HCl, 0.1 M Na₂HPO₄, pH 8.0 for 1 hour with vigorous shaking. The recombinant *Dac g 1* was purified from the extract on a Ni⁺² chelating column (Hochuli et al. (1987) *J. Chromatog.*, **411**:177-184; Hochuli et al. (1988), *Bio/Tech.*, **6**:1321-1325).

30 B. Cloning of *Poa p I*.

- RNA was isolated from the pollen of *Poa pratensis*, double stranded cDNA was prepared and self-annealed oligonucleotides AT and AL were added as described in section A, above. PCR product was amplified using oligonucleotide primers PhI-7 (5'-CCGAATTCGTGGAGAAGGGGTCCAA-3') (SEQ ID NO: 90) and Poa-1 (5'-
 35 TTAGGATCCTCACTTATCATAIGACGTATC-3' (SEQ ID NO: 91), wherein C at

position 13 can also be T, A at position 16 can also be G, A at position 19 can also be G, G at position 23 can also be C, A at position 24 can also be T, C at position 25 can also be T or A or G and A at position 28 can be G). All *Poa p 1* clones were amplified by 20 cycles of heating to 94°C for 1 min, 55°C for 1 min and 72°C for 1 min. The amplified material was finally heated to 72°C for 5 min. Three clones, 11, 15 and 17, were isolated that contained part of the nucleotide sequence for the gene that encodes *Poa p 1*. The *Dac g 1* sequence encoded by clones 11, 15 and 17 corresponds to amino acids 151 - 240 of Fig. 6.

Clones containing partial nucleotide sequences of the gene encoding *Poa p 1* were derived from PCRs that used oligonucleotide primers AP and Poa-3 (5'-TTGAATTCCTTGTCATTGCCCTTCTG-3') (SEQ ID NO: 92) in the primary PCR and AP and Poa-4 (5'-AAGAATTCCTTCTGCTTGATGTCCAC-3') (SEQ ID NO: 93) in the secondary PCR. Other clones were derived from PCRs that used oligonucleotide primers AP and Poa-6 (5'-ATGAATTCGAGTCGTGGGGAGCCGTC-3') (SEQ ID NO: 94) in the primary PCR and AP and Poa-7 (5'-ATGAATTCGTCTGGAGGATCGACACC-3') (SEQ ID NO: 95) in the secondary PCR. Clones 58, 59 and 63 were derived from the PCR using primers AP and Poa-4. Clones 91 and 97 were derived from the PCR using primers AP and Poa-7.

Additional clones were derived from a PCR that used oligonucleotide primers Poa-1 and Poa-5 (5'-ATGAATTCATCGCAAAGGTTCCCCC-3' (SEQ ID NO: 96), wherein A at position 14 can also be G or C or T). These clones, 113, 114 and 115, corresponded to the portion of the gene that encoded amino acids 1 - 240 of *Poa p 1* (see Fig. 6). The nucleotide (SEQ ID NO: 53) and deduced amino acid (SEQ ID NO: 54) sequences of clone 114 are shown in Fig. 6. Nucleotide 93 in Fig. 6 was not resolved and could be a G or a C or a T or an A and is represented by the letter "N". Nucleotide 94 in Fig. 6 was not conclusively resolved and could be a G or a C or a T but not an A and is represented by the letter "B". The codon containing nucleotide 93 (GGN) encodes a Glycine at residue 31. The codon containing nucleotide 94 (BCC) encodes an Alanine (GCC), a Proline (CCC), or a Serine (TCC) at amino acid 32. The amino acid at residue 32 in Fig. 6 is represented by an "X".

Inserts from clones 11 and 114 were isolated and ligated into appropriately digested expression vector pET-11d (Novagen, Madison, WI: Jameel et al. (1990) *J. Virol.* 64:3963-3966). Recombinant proteins were expressed as described in section A, above.

C. Cloning of *Phl p 1*.

RNA was isolated from the pollen of *Phleum pratense*, double stranded cDNA was prepared and self-annealed oligonucleotides AT and AL were added as described in section A, above. Clones were derived from a PCR that used oligonucleotide primers PhA1.1 (5'-TTTGGATCCTCACTTGGACTCGTAGCT-3') (SEQ ID NO: 97) and Phl-2 (5'-TTGAATTCTCGCGAAGGTGCCCCCG-3') (SEQ ID NO: 98), wherein G at position 13 can also be A). These clones, 20 and 22, corresponded to the portion of the gene that encoded amino acids 1 - 240 of *Phl p 1* (see Fig. 7). The nucleotide (SEQ ID NO: 55) and deduced amino acid (SEQ ID NO: 56) sequences of clone 20 are shown in Fig. 7.

Clones containing partial nucleotide sequence of the gene encoding *Phl p 1* were derived from a PCR using oligonucleotide primers Phl-7 (5'-CCGAATTCGTGGAGAAGGGGTCCAA-3') (SEQ ID NO: 90) and PhA1.1. Clones 47-52 were derived from this PCR. These clones encoded amino acids 151 through 240 of Fig. 7.

Inserts from clones 22 and 51 were isolated and ligated into appropriately digested expression vector pET-11d (Novagen, Madison, WI: Jameel et al. (1990) *J. Virol.* 64:3963-3966). Recombinant proteins were expressed as described in section A, above.

Example 6 - Comparison of *Dac g 1*, *Phl p 1* and *Poa p 1* With *Lol p 1*.

The sequences for *Dac g 1* (Fig. 5) (SEQ ID NO: 58), *Phl p 1* (Fig. 7) (SEQ ID NO: 59) and *Poa p 1* (Fig. 6) (SEQ ID NO: 60) were compared with *Lol p 1* (SEQ ID NO: 57). The amino acid sequences of these Group 1 allergens had 95% (*Dac g 1*), 91% (*Phl p 1*) and 91% (*Poa p 1*) identity, respectively, with *Lol p 1*. This comparison is shown schematically in Fig. 8. The complete sequence of *Lol p 1* is shown in standard one letter code. Only differences from the *Lol p 1* sequence are shown for the other Group 1 allergens; identity is indicated by a dash (-). Potential amino acid polymorphisms were predicted by detected nucleotide polymorphisms in each sequence. Such potential polymorphisms are shown by superscript and subscript letters at the site of the polymorphism.

T cell epitope containing peptides of *Lol p 1*, peptides 16.1 (SEQ ID NO: 23), 18 (SEQ ID NO: 25), 20 (SEQ ID NO: 27) and 23 (SEQ ID NO: 30), were defined in

Example 4 (Fig. 3). The sequences of the other Group 1 allergens are very conserved in these regions. Since the Group 1 allergens are homologous, the major T cell epitope containing peptides of *Lol p 1* are likely to be the major T cell epitope containing regions in the related grasses. Comparison of the sequences of the *Lol p 1* peptides with the homologous peptides containing *Dac g 1*, *Phl p 1* and *Poa p 1* polymorphisms are shown in Fig. 9 (SEQ ID NO: 23, 25, 27, 30, 61-70).

SEQUENCE LISTING

- (1) GENERAL INFORMATION:
- 5 (i) APPLICANT:
- (A) NAME: IMMULOGIC PHARMACEUTICAL CORPORATION
 - (B) STREET: 610 LINCOLN STREET
 - (C) CITY: WALTHAM
 - 10 (D) STATE: MASSACHUSETTS
 - (E) COUNTRY: USA
 - (F) POSTAL CODE (ZIP): 02154
 - (G) TELEPHONE: (617) 466-6000
 - 15 (H) TELEFAX: (617) 466-6010
- (ii) TITLE OF INVENTION: T CELL EPITOPES OF RYEGRASS POLLEN ALLERGENS
- (iii) NUMBER OF SEQUENCES: 98
- 20 (iv) CORRESPONDENCE ADDRESS:
- (A) ADDRESSEE: LAHIVE & COCKFIELD
 - (B) STREET: 60 State Street, suite # 510
 - (C) CITY: Boston
 - 25 (D) STATE: Massachusetts
 - (E) COUNTRY: US
 - (F) ZIP: 02109-1875
- (v) COMPUTER READABLE FORM:
- 30 (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: ASCII text
- 35 (vi) CURRENT APPLICATION DATA:
- (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- 40 (vii) PRIOR APPLICATION DATA:
- (A) APPLICATION NUMBER: US 08/106,016
 - (B) FILING DATE: 31-AUG-1993
- (vii) PRIOR APPLICATION DATA:
- 45 (A) APPLICATION NUMBER: US 08/031,001
 - (B) FILING DATE: 12-MAR-1993
- (viii) ATTORNEY/AGENT INFORMATION:
- 50 (A) NAME: Amy E. Mandragouras
 - (B) REGISTRATION NUMBER: 36,207
 - (C) REFERENCE/DOCKET NUMBER: (IMI-040PC)
- (ix) TELECOMMUNICATION INFORMATION:
- 55 (A) TELEPHONE: (617) 227-7400
 - (B) TELEFAX: (617) 227-5941
- (2) INFORMATION FOR SEQ ID NO:1:
- 60 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1124 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

5 (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 16..804

10 (ix) FEATURE:
 (A) NAME/KEY: mat_peptide
 (B) LOCATION: 85..804

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

15 CAAATTCAAG ACAAG ATG GCG TCC TCC TCG TCG GTG CTC CTG GTG GTG GCG
 51
 Met Ala Ser Ser Ser Ser Val Leu Leu Val Val Ala
 -23 -20 -15

20 CTG TTC GCC GTG TTC CTG GGC AGC GCG CAT GGC ATC GCG AAG GTA CCA
 99
 Leu Phe Ala Val Phe Leu Gly Ser Ala His Gly Ile Ala Lys Val Pro
 -10 -5 1 5

25 CCG GGC CCC AAC ATC ACG GCC GAG TAC GGC GAC AAG TGG CTG GAC GCG
 147
 Pro Gly Pro Asn Ile Thr Ala Glu Tyr Gly Asp Lys Trp Leu Asp Ala
 10 15 20

30 AAG AGC ACC TGG TAT GGC AAG CCG ACC GGC GCC GGT CCC AAG GAC AAC
 195
 Lys Ser Thr Trp Tyr Gly Lys Pro Thr Gly Ala Gly Pro Lys Asp Asn
 25 30 35

35 GGC GGC GCG TGC GGG TAC AAG GAC GTT GAC AAG GCG CCG TTC AAC GGC
 243
 Gly Gly Ala Cys Gly Tyr Lys Asp Val Asp Lys Ala Pro Phe Asn Gly
 40 45 50

40 ATG ACC GGC TGC GGC AAC ACC CCC ATC TTC AAG GAC GGC CGT GGC TGC
 291
 Met Thr Gly Cys Gly Asn Thr Pro Ile Phe Lys Asp Gly Arg Gly Cys
 55 60 65

45 GGC TCC TGC TTC GAG ATC AAG TGC ACC AAG CCC GAG TCC TGC TCC GGC
 339
 Gly Ser Cys Phe Glu Ile Lys Cys Thr Lys Pro Glu Ser Cys Ser Gly
 70 75 80 85

50 GAG GCT GTC ACC GTC ACA ATC ACC GAC GAC AAC GAG GAG CCC ATC GCA
 387
 Glu Ala Val Thr Val Thr Ile Thr Asp Asp Asn Glu Glu Pro Ile Ala
 90 95 100

55 CCC TAC CAT TTC GAC CTC TCG GGC CAC GCG TTC GGG TCC ATG GCG AAG
 435
 Pro Tyr His Phe Asp Leu Ser Gly His Ala Phe Gly Ser Met Ala Lys
 105 110 115

60 AAG GGC GAG GAG CAG AAG CTC CGC AGC GCC GGC GAG CTG GAG CTC CAG
 483
 Lys Gly Glu Glu Gln Lys Leu Arg Ser Ala Gly Glu Leu Glu Leu Gln
 120 125 130

5 TTC AGG CGG GTC AAG TGC AAG TAC CCG GAC GGC ACC AAG CCG ACA TTC
 531
 Phe Arg Arg Val Lys Cys Lys Tyr Pro Asp Gly Thr Lys Pro Thr Phe
 135 140 145

 10 CAC GTC GAG AAG GCT TCC AAC CCC AAC TAC CTC GCT ATT CTG GTG AAG
 579
 His Val Glu Lys Ala Ser Asn Pro Asn Tyr Leu Ala Ile Leu Val Lys
 150 155 160 165

 15 TAC GTC GAC GGC GAC GGT GAC GTG GTG GCG GTG GAC ATC AAG GAG AAG
 627
 Tyr Val Asp Gly Asp Gly Asp Val Val Ala Val Asp Ile Lys Glu Lys
 170 175 180

 20 GGC AAG GAT AAG TGG ATC GAG CTC AAG GAG TCG TGG GGA GCA GTC TGG
 675
 Gly Lys Asp Lys Trp Ile Glu Leu Lys Glu Ser Trp Gly Ala Val Trp
 185 190 195

 25 AGG ATC GAC ACC CCC GAT AAG CTG ACG GGC CCA TTC ACC GTC CGC TAC
 723
 Arg Ile Asp Thr Pro Asp Lys Leu Thr Gly Pro Phe Thr Val Arg Tyr
 200 205 210

 30 ACC ACC GAG GGC GGC ACC AAA TCC GAA GTC GAG GAT GTC ATC CCT GAG
 771
 Thr Thr Glu Gly Gly Thr Lys Ser Glu Val Glu Asp Val Ile Pro Glu
 215 220 225

 35 GGC TGG AAG GCC GAC ACC TCC TAC TCG GCC AAG TGAGCAAGAA GTGGAGTGAT
 824
 Gly Trp Lys Ala Asp Thr Ser Tyr Ser Ala Lys
 230 235 240

 CTTCTTCCAA TCAGCTTAAT TTTGACTCAA GATCTCAAAT AATCCAGCCG CACATATATA
 884

 40 CGAGGCGGTG AGACATACAA GCTCCTCCAT GAGTATATTC ATTCATGCCG TATAGAGAGG
 944

 AGAAAGATGC CTGAATAAGA GTTTGAGGTC GACACCTTGT GAGAAGTGTA TATAGGAGGA
 1004

 45 ACCCAATCTG GCTCCATCTT TCTTTGCTCG CACGGTGATC TGCTAAGGTT ATCTTCTAAC
 1064

 50 AGGCCAGATT AACCTACTAT CTAATATATG CAACGTATGG TCATTTTCCC TAAAAA
 1124

(2) INFORMATION FOR SEQ ID NO:2:

55 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 263 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

60 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Ala Ser Ser Ser Ser Val Leu Leu Val Val Ala Leu Phe Ala Val

	-23		-20				-15				-10			
	Phe	Leu	Gly	Ser	Ala	His	Gly	Ile	Ala	Lys	Val	Pro	Pro	Asn
			-5					1				5		
5	Ile	Thr	Ala	Glu	Tyr	Gly	Asp	Lys	Trp	Leu	Asp	Ala	Lys	Trp
	10					15					20			25
10	Tyr	Gly	Lys	Pro	Thr	Gly	Ala	Gly	Pro	Lys	Asp	Asn	Gly	Cys
					30					35			40	
	Gly	Tyr	Lys	Asp	Val	Asp	Lys	Ala	Pro	Phe	Asn	Gly	Met	Cys
				45					50			55		
15	Gly	Asn	Thr	Pro	Ile	Phe	Lys	Asp	Gly	Arg	Gly	Cys	Gly	Phe
			60					65				70		
	Glu	Ile	Lys	Cys	Thr	Lys	Pro	Glu	Ser	Cys	Ser	Gly	Glu	Thr
	75						80					85		
20	Val	Thr	Ile	Thr	Asp	Asp	Asn	Glu	Glu	Pro	Ile	Ala	Pro	Phe
	90					95					100			105
25	Asp	Leu	Ser	Gly	His	Ala	Phe	Gly	Ser	Met	Ala	Lys	Lys	Glu
					110					115				120
	Gln	Lys	Leu	Arg	Ser	Ala	Gly	Glu	Leu	Glu	Leu	Gln	Phe	Val
				125					130				135	
30	Lys	Cys	Lys	Tyr	Pro	Asp	Gly	Thr	Lys	Pro	Thr	Phe	His	Lys
			140					145					150	
	Ala	Ser	Asn	Pro	Asn	Tyr	Leu	Ala	Ile	Leu	Val	Lys	Tyr	Gly
	155						160					165		
35	Asp	Gly	Asp	Val	Val	Ala	Val	Asp	Ile	Lys	Glu	Lys	Gly	Lys
	170					175					180			185
40	Trp	Ile	Glu	Leu	Lys	Glu	Ser	Trp	Gly	Ala	Val	Trp	Arg	Thr
					190					195				200
	Pro	Asp	Lys	Leu	Thr	Gly	Pro	Phe	Thr	Val	Arg	Tyr	Thr	Glu
				205					210				215	
45	Gly	Thr	Lys	Ser	Glu	Val	Glu	Asp	Val	Ile	Pro	Glu	Gly	Ala
			220					225					230	
50	Asp	Thr	Ser	Tyr	Ser	Ala	Lys							
	235						240							

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 20 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- 10 (v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

15 Ile Ala Lys Val Pro Pro Gly Pro Asn Ile Thr Ala Glu Tyr Gly Asp
1 5 10 15
20 Lys Trp Leu Asp
20

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 20 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- 30 (v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

35 Ile Ala Lys Val Xaa Pro Gly Xaa Asn Ile Thr Ala Glu Tyr Gly Asp
1 5 10 15
40 Lys Trp Leu Asp
20

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- 45 (A) LENGTH: 20 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- 50 (v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

55 Thr Ala Glu Tyr Gly Asp Lys Trp Leu Asp Ala Lys Ser Thr Trp Tyr
1 5 10 15
60 Gly Lys Pro Thr
20

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: peptide
 (v) FRAGMENT TYPE: internal

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

15 Ala Lys Ser Thr Trp Tyr Gly Lys Pro Thr Gly Ala Gly Pro Lys Asp
 1 5 10 15
 Asn Gly Gly Ala
 20

20 (2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 25 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
 (v) FRAGMENT TYPE: internal

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

35 Gly Ala Gly Pro Lys Asp Asn Gly Gly Ala Cys Gly Tyr Lys Asn Val
 1 5 10 15
 Asp Lys Ala Pro
 20

40 (2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 45 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide
 (v) FRAGMENT TYPE: internal

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

55 Gly Ala Gly Pro Lys Asp Asn Gly Gly Ala Cys Gly Tyr Lys Asp Val
 1 5 10 15
 Asp Lys Ala Pro
 20

60 (2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: peptide
(v) FRAGMENT TYPE: internal

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Cys Gly Tyr Lys Asp Val Asp Lys Ala Pro Phe Asn Gly Met Thr Gly
1 5 10 15

15

Cys Gly Asn Thr
20

20 (2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: peptide
(v) FRAGMENT TYPE: internal

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Phe Asn Gly Met Thr Gly Cys Gly Asn Thr Pro Ile Phe Lys Asp Gly
1 5 10 15

35

Arg Gly Cys Gly
20

5 (2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

20 Pro Ile Phe Lys Asp Gly Arg Gly Cys Gly Ser Cys Phe Glu Ile Lys
1 5 10 15

Cys Thr Lys Pro
20

25 (2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

40 Ser Cys Phe Glu Ile Lys Cys Thr Lys Pro Glu Ser Cys Ser Gly Glu
1 5 10 15

Ala Val Thr Val
20

45

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

60 Glu Ser Cys Ser Gly Glu Ala Val Thr Val Thr Ile Thr Asp Asp Asn
1 5 10 15

Glu Glu Pro Ile

20

(2) INFORMATION FOR SEQ ID NO:14:

- 5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Thr Ile Thr Asp Asp Asn Glu Glu Pro Ile Ala Pro Tyr His Phe Asp
 1 5 10 15
 Leu Ser Gly His
 20

20

(2) INFORMATION FOR SEQ ID NO:15:

25

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Ala Pro Tyr His Phe Asp Leu Ser Gly His Ala Phe Gly Ser Met Ala
 1 5 10 15
 Asp Asp Gly Glu
 20

40

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 20 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- 10 (v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

- 15 Ala Pro Tyr His Phe Asp Leu Ser Gly His Ala Phe Gly Ser Met Ala
1 5 10 15
20 Lys Lys Gly Glu
20

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 20 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- 30 (v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

- 35 Ala Phe Gly Ser Met Ala Asp Asp Gly Glu Glu Gln Lys Leu Arg Ser
1 5 10 15
40 Ala Gly Glu Leu
20

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- 45 (A) LENGTH: 20 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- 50 (v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

- 55 Ala Phe Gly Ser Met Ala Lys Lys Gly Glu Glu Gln Lys Leu Arg Ser
1 5 10 15
60 Ala Gly Glu Leu
20

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Glu Gln Lys Leu Arg Ser Ala Gly Glu Leu Glu Leu Gln Phe Arg Arg
 1 5 10 15
 Val Lys Cys Lys
 20

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Glu Leu Gln Phe Arg Arg Val Lys Cys Lys Tyr Pro Asp Asp Thr Lys
 1 5 10 15
 Pro Thr Phe His
 20

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Tyr Pro Asp Asp Thr Lys Pro Thr Phe His Val Glu Lys Ala Ser Asn
 1 5 10 15
 Pro Asn Tyr Leu
 20

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
5
(ii) MOLECULE TYPE: peptide
(v) FRAGMENT TYPE: internal
10
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:
Val Glu Lys Ala Ser Asn Pro Asn Tyr Leu Ala Ile Leu Val Lys Tyr
15 1 5 10 15
Val Asp Gly Asp
20
(2) INFORMATION FOR SEQ ID NO:23:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
25 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: peptide
(v) FRAGMENT TYPE: internal
30
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:
Val Glu Lys Gly Ser Asn Pro Asn Tyr Leu Ala Ile Leu Val Lys Tyr
35 1 5 10 15
Val Asp Gly Asp
40 20
(2) INFORMATION FOR SEQ ID NO:24:
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
45 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: peptide
50 (v) FRAGMENT TYPE: internal
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:
55 Ala Ile Leu Val Lys Tyr Val Asp Gly Asp Gly Asp Val Val Ala Val
1 5 10 15
Asp Ile Lys Glu
60 20
(2) INFORMATION FOR SEQ ID NO:25:
(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: peptide
 (v) FRAGMENT TYPE: internal

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

Gly Asp Val Val Ala Val Asp Ile Lys Glu Lys Gly Lys Asp Lys Trp
 1 5 10 15

15 Ile Glu Leu Lys
 20

(2) INFORMATION FOR SEQ ID NO:26:

20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: peptide
 (v) FRAGMENT TYPE: internal

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

Lys Gly Lys Asp Lys Trp Ile Glu Leu Lys Glu Ser Trp Gly Ala Val
 1 5 10 15

Trp Arg Ile Asp
 20

40 (2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: peptide
 (v) FRAGMENT TYPE: internal

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

Glu Ser Trp Gly Ala Val Trp Arg Ile Asp Thr Pro Asp Lys Leu Thr
 1 5 10 15

Gly Pro Phe Thr
 20

60

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids

(B) TYPE: amino acid
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: peptide
(v) FRAGMENT TYPE: internal

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Thr	Pro	Asp	Lys	Leu	Thr	Gly	Pro	Phe	Thr	Val	Arg	Tyr	Thr	Thr	Glu
1				5					10					15	

15 Gly Gly Thr Lys
20

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

5

- (A) LENGTH: 20 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

10

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

15

Val Arg Tyr Thr Thr Glu Gly Gly Thr Lys Ser Glu Val Glu Asp Val
 1 5 10 15

20

Ile Pro Glu Gly
 20

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

25

- (A) LENGTH: 20 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

30

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

35

Ser Glu Val Glu Asp Val Ile Pro Glu Gly Trp Lys Ala Asp Thr Ser
 1 5 10 15

40

Tyr Ser Ala Lys
 20

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

45

- (A) LENGTH: 22 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

Asp Glu Val Glu Lys Gly Ser Asn Pro Asn Tyr Leu Ala Ile Leu Val
 1 5 10 15

60

Lys Tyr Val Asp Gly Asp
 20

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 amino acids
 (B) TYPE: amino acid
5 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

15 Asp Glu Ala Glu Lys Gly Ser Asn Pro Asn Tyr Leu Ala Ile Leu Val
 1 5 10 15

 Lys Tyr Val Asp Gly Asp
 20

20

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 amino acids
25 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

30 (v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

35 Lys Lys Val Glu Lys Gly Ser Asn Pro Asn Tyr Leu Ala Ile Leu Val
 1 5 10 15

 Lys Lys

40

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:
45 (A) LENGTH: 15 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

50 (v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

55 Val Glu Lys Gly Ser Asn Pro Asn Tyr Leu Ala Ile Leu Asp Glu
 1 5 10 15

60

(2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 amino acids
 (B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

5 (v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

10 Ala Glu Lys Gly Ser Asn Pro Asn Tyr Leu Ala Ile Leu Asp Glu
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:36:

15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 16 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

30 Asp Glu Val Glu Lys Gly Ser Asn Pro Asn Tyr Leu Ala Ile Asp Glu
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:37:

35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

Lys Lys Ala Glu Lys Gly Ser Asn Pro Asn Tyr Leu Ala Ile Leu Val
1 5 10 15

50 Lys Lys

(2) INFORMATION FOR SEQ ID NO:38:

55 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

60 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

5 Asp Glu Pro Asn Tyr Leu Ala Ile Leu Val Lys Tyr Val Asp Glu
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:39:

10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 15 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

```

15      (ii) MOLECULE TYPE: peptide
      (v) FRAGMENT TYPE: internal

```

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

25 Gly Asp Val Val Ala Val Asp Ile Lys Glu Lys Gly Lys Asp Lys
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:

5 (A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

10 (v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

15 Val Ala Val Asp Ile Lys Glu Lys Gly Lys Asp Lys Trp Ile Glu
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:41:

20 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

35 Ala Val Asp Ile Lys Glu Lys Gly Lys Asp Lys Trp Ile Glu Leu
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:42:

40 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 14 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

Asp Ile Lys Glu Lys Gly Lys Asp Lys Trp Ile Glu Leu Lys
1 5 10

(2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 14 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- 10 (v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

- 15 Trp Gly Ala Val Trp Arg Ile Asp Thr Pro Asp Lys Leu Thr
1 5 10

(2) INFORMATION FOR SEQ ID NO:44:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 14 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

- 25 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

- 35 Gly Ala Val Trp Arg Ile Asp Thr Pro Asp Lys Leu Thr Gly
1 5 10

(2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:

- 40 (A) LENGTH: 14 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- 45 (v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

- 50 Trp Arg Ile Asp Thr Pro Asp Lys Leu Thr Gly Pro Phe Thr
1 5 10

(2) INFORMATION FOR SEQ ID NO:46:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 14 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

10 (v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

15 Glu Ser Trp Gly Ala Val Trp Arg Ile Asp Thr Pro Asp Lys
1 5 10

(2) INFORMATION FOR SEQ ID NO:47:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 14 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

35 Ala Gly Ala Val Trp Arg Ile Asp Thr Pro Asp Lys Leu Thr
1 5 10

(2) INFORMATION FOR SEQ ID NO:48:

(i) SEQUENCE CHARACTERISTICS:

- 40 (A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

45 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

Ser Glu Val Glu Asp Val Ile Pro Glu Gly Trp Lys Ala Asp Thr
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:49:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- 10 (v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

15 Glu Asp Val Ile Pro Glu Gly Trp Lys Ala Asp Thr Ser Tyr Ser
1 5 10 15

(2) INFORMATION FOR SEQ ID NO:50:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 14 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

35 Ile Pro Glu Gly Trp Lys Ala Asp Thr Ser Tyr Ser Ala Lys
1 5 10

(2) INFORMATION FOR SEQ ID NO:51:

(i) SEQUENCE CHARACTERISTICS:

- 40 (A) LENGTH: 723 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

45

(ix) FEATURE:

- 50 (A) NAME/KEY: CDS
(B) LOCATION: 1..720

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

	ATC	CCG	AAG	GTG	CCC	CCG	GGC	CCG	AAC	ATC	ACG	GCG	ACC	TAC	GGT	GAC	48
5	Ile	Pro	Lys	Val	Pro	Pro	Gly	Pro	Asn	Ile	Thr	Ala	Thr	Tyr	Gly	Asp	15
	1				5					10							
	AAG	TGG	CTG	GAC	GCG	AAG	AGC	ACA	TGG	TAC	GGC	AAG	CCG	ACG	GGC	GCC	96
10	Lys	Trp	Leu	Asp	Ala	Lys	Ser	Thr	Trp	Tyr	Gly	Lys	Pro	Thr	Gly	Ala	30
				20					25								
	GGC	CCC	AAG	GAC	AAC	GGC	GGC	GCG	TGC	GGG	TAC	AAG	GAC	GTG	GAC	AAG	144
15	Gly	Pro	Lys	Asp	Asn	Gly	Gly	Ala	Cys	Gly	Tyr	Lys	Asp	Val	Asp	Lys	45
			35					40									
	GCG	CCG	TTC	AAC	GGC	ATG	ACC	GGG	TGC	GGC	AAC	ACC	CCC	ATC	TTC	AAG	192
20	Ala	Pro	Phe	Asn	Gly	Met	Thr	Gly	Cys	Gly	Asn	Thr	Pro	Ile	Phe	Lys	55
		50					55					60					
	GAC	GGG	CGC	GGG	TGC	GGT	TCC	TGC	TTC	GAG	ATC	AAG	TGC	ACG	AAG	CCC	240
25	Asp	Gly	Arg	Gly	Cys	Gly	Ser	Cys	Phe	Glu	Ile	Lys	Cys	Thr	Lys	Pro	70
	65					70					75					80	
	GAG	TCG	TGC	TCC	GGC	GAG	GCC	GTC	ACC	GTC	CAC	ATC	ACC	GAC	GAC	AAC	288
30	Glu	Ser	Cys	Ser	Gly	Glu	Ala	Val	Thr	Val	His	Ile	Thr	Asp	Asp	Asn	85
					85					90					95		
	GAG	GAG	CCC	ATC	GCG	CCC	TAC	CAC	TTC	GAC	CTT	TCC	GGC	CAC	GCG	TTC	336
35	Glu	Glu	Pro	Ile	Ala	Pro	Tyr	His	Phe	Asp	Leu	Ser	Gly	His	Ala	Phe	100
				100					105					110			
	GGT	TCC	ATG	GCG	AAG	AAG	GGC	GAG	GAG	CAG	AAG	CTG	CGC	AGC	GCG	GGC	384
40	Gly	Ser	Met	Ala	Lys	Lys	Gly	Glu	Glu	Gln	Lys	Leu	Arg	Ser	Ala	Gly	115
			115				120					125					
	GAG	CTG	GAG	CTG	CAG	TTT	AGG	CGG	GTG	AAG	TGC	AAG	TAC	CCC	GAG	GGC	432
45	Glu	Leu	Glu	Leu	Gln	Phe	Arg	Arg	Val	Lys	Cys	Lys	Tyr	Pro	Glu	Gly	130
		130				135						140					
	ACC	AAG	GTG	ACC	TTC	CAC	GTC	GAG	AAG	GGT	TCC	AAC	CCC	AAC	TAC	CTG	480
50	Thr	Lys	Val	Thr	Phe	His	Val	Glu	Lys	Gly	Ser	Asn	Pro	Asn	Tyr	Leu	145
					150						155					160	
	GCG	CTG	CTG	GTG	AAG	TAC	GTC	GAC	GGC	GAC	GGC	GAC	GTG	GTG	GCG	GTG	528
55	Ala	Leu	Leu	Val	Lys	Tyr	Val	Asp	Gly	Asp	Gly	Asp	Val	Val	Ala	Val	165
					165					170					175		
	GAT	ATC	AAG	GAG	AAG	GGC	AAG	GAC	AAG	TGG	ATC	GCG	CTC	AAG	GAG	TCA	576
60	Asp	Ile	Lys	Glu	Lys	Gly	Lys	Asp	Lys	Trp	Ile	Ala	Leu	Lys	Glu	Ser	180
				180					185					190			
	TGG	GGA	GCC	ATC	TGG	AGG	GTG	GAC	ACC	CCC	GAC	AAG	CTG	ACG	GGC	CCA	624

Trp	Gly	Ala	Ile	Trp	Arg	Val	Asp	Thr	Pro	Asp	Lys	Leu	Thr	Gly	Pro
		195					200					205			

TTC ACC GTT CGC TAC ACC ACC GAG GGA GGC ACC AAG TCC GAA GTT GAG
 672
 Phe Thr Val Arg Tyr Thr Thr Glu Gly Gly Thr Lys Ser Glu Val Glu
 210 215 220
 5
 GAC GTC ATC CCC GAG GGC TGG AAG GCC GAC GCC AGC TAC GAG TCC AAG
 720
 Asp Val Ile Pro Glu Gly Trp Lys Ala Asp Ala Ser Tyr Glu Ser Lys
 225 230 235 240
 10
 TGA
 723

(2) INFORMATION FOR SEQ ID NO:52:

15 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 240 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: protein

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

Ile Pro Lys Val Pro Pro Gly Pro Asn Ile Thr Ala Thr Tyr Gly Asp
 1 5 10 15
 30 Lys Trp Leu Asp Ala Lys Ser Thr Trp Tyr Gly Lys Pro Thr Gly Ala
 20 25 30
 Gly Pro Lys Asp Asn Gly Gly Ala Cys Gly Tyr Lys Asp Val Asp Lys
 35 35 40 45
 Ala Pro Phe Asn Gly Met Thr Gly Cys Gly Asn Thr Pro Ile Phe Lys
 50 55 60
 40 Asp Gly Arg Gly Cys Gly Ser Cys Phe Glu Ile Lys Cys Thr Lys Pro
 65 70 75 80
 Glu Ser Cys Ser Gly Glu Ala Val Thr Val His Ile Thr Asp Asp Asn
 85 90 95
 45 Glu Glu Pro Ile Ala Pro Tyr His Phe Asp Leu Ser Gly His Ala Phe
 100 105 110
 Gly Ser Met Ala Lys Lys Gly Glu Glu Gln Lys Leu Arg Ser Ala Gly
 115 120 125
 50 Glu Leu Glu Leu Gln Phe Arg Arg Val Lys Cys Lys Tyr Pro Glu Gly
 130 135 140
 55 Thr Lys Val Thr Phe His Val Glu Lys Gly Ser Asn Pro Asn Tyr Leu
 145 150 155 160
 Ala Leu Leu Val Lys Tyr Val Asp Gly Asp Gly Asp Val Val Ala Val
 165 170 175
 60 Asp Ile Lys Glu Lys Gly Lys Asp Lys Trp Ile Ala Leu Lys Glu Ser
 180 185 190
 Trp Gly Ala Ile Trp Arg Val Asp Thr Pro Asp Lys Leu Thr Gly Pro
 195 200 205

Phe Thr Val Arg Tyr Thr Thr Glu Gly Gly Thr Lys Ser Glu Val Glu
210 215 220

5 Asp Val Ile Pro Glu Gly Trp Lys Ala Asp Ala Ser Tyr Glu Ser Lys
225 230 235 240

(2) INFORMATION FOR SEQ ID NO:53:

10 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 723 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: cDNA

20 (ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 1..720

(ix) FEATURE:
(A) NAME/KEY: Modified-site
25 (B) LOCATION: 32
(D) OTHER INFORMATION: /note= "Xaa is Ser, Pro or Ala"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

30 ATC GCG AAG GTT CCC CCC GGC CCG AAC ATC ACG GCG ACC TAC GGC GAC
48
Ile Ala Lys Val Pro Pro Gly Pro Asn Ile Thr Ala Thr Tyr Gly Asp
1 5 10 15
35 AAG TGG CTT GAC GCG AAG AGC ACC TGG TAC GGC AAG CCG ACC GGN BCC
96
Lys Trp Leu Asp Ala Lys Ser Thr Trp Tyr Gly Lys Pro Thr Gly Xaa
20 25 30
40 GGT CCC AAG GAC AAC GGC GGC GCG TGC GGA TAC AAG GAC GTG GAC AAG
144
Gly Pro Lys Asp Asn Gly Gly Ala Cys Gly Tyr Lys Asp Val Asp Lys
35 40 45
45 CCC CCG TTC AGC GGC ATG ACC GGC TGC GGC AAC ACC CCC ATC TTC AAG
192
Pro Pro Phe Ser Gly Met Thr Gly Cys Gly Asn Thr Pro Ile Phe Lys
50 55 60
50 TCC GGC CGC GGC TGC GGC TCC TGC TTC GAG ATC AAG TGC ACC AAG CCC
240
Ser Gly Arg Gly Cys Gly Ser Cys Phe Glu Ile Lys Cys Thr Lys Pro
65 70 75 80
55 GAG TCC TGC TCC GGG GAG CCC GTC CTG GTC CAC ATC ACC GAC GAC AAC
288
Glu Ser Cys Ser Gly Glu Pro Val Leu Val His Ile Thr Asp Asp Asn
85 90 95
60 GAG GAG CCC ATC GCC GCC TAC CAC TTC GAC CTC TCC GGC AAG GCG TTC
336
Glu Glu Pro Ile Ala Ala Tyr His Phe Asp Leu Ser Gly Lys Ala Phe
100 105 110

5 GGG GCC ATG GCC AAG AAG GGT GAG GAG CAG AAG CTG CGC AGC GCC GGC
 384
 Gly Ala Met Ala Lys Lys Gly Glu Glu Gln Lys Leu Arg Ser Ala Gly
 115 120 125
 10 GAG CTG GAG CTC AAG TTC CGC CGC GTC AAG TGC GAG TAC CCG AAG GGC
 432
 Glu Leu Glu Leu Lys Phe Arg Arg Val Lys Cys Glu Tyr Pro Lys Gly
 130 135 140
 15 ACC AAG GTT ACC TTC CAC GTC GAG AAG GGG TCC AAC CCC AAC TAC CTT
 480
 Thr Lys Val Thr Phe His Val Glu Lys Gly Ser Asn Pro Asn Tyr Leu
 145 150 155 160
 20 GCG CTG CTG GTG AAG TAC GTC GAC GGC GAC GGG GAC GTG GTG GCG GTG
 528
 Ala Leu Leu Val Lys Tyr Val Asp Gly Asp Gly Asp Val Val Ala Val
 165 170 175
 25 GAC ATC AAG CAG AAG GGC AAG GAC AAG TGG ATC GAG CTC AAG GAG TCG
 576
 Asp Ile Lys Gln Lys Gly Lys Asp Lys Trp Ile Glu Leu Lys Glu Ser
 180 185 190
 30 TGG GGA GCC GTC TGG AGG ATC GAC ACC CCC GAC AAG CTC ACC GGC CCC
 624
 Trp Gly Ala Val Trp Arg Ile Asp Thr Pro Asp Lys Leu Thr Gly Pro
 195 200 205
 35 TTC ACC GTC CGC TAC ACC ACC GAG GGC GGC ACC AAG GCC GAA GCC GAG
 672
 Phe Thr Val Arg Tyr Thr Thr Glu Gly Gly Thr Lys Ala Glu Ala Glu
 210 215 220
 40 GAC GTC ATC CCC GAG GGC TGG AAG GCC GAC ACC GCC TAC GAG GCC AAG
 720
 Asp Val Ile Pro Glu Gly Trp Lys Ala Asp Thr Ala Tyr Glu Ala Lys
 225 230 235 240
 TGA
 723

45 (2) INFORMATION FOR SEQ ID NO:54:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 240 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

50

(ii) MOLECULE TYPE: protein

55

(ix) FEATURE:

(A) NAME/KEY: Modified-site

(B) LOCATION: 32

(D) OTHER INFORMATION: /note= "Xaa is Ser, Pro or Ala"

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

```

Ile Ala Lys Val Pro Pro Gly Pro Asn Ile Thr Ala Thr Tyr Gly Asp
 1          5          10          15
10 Lys Trp Leu Asp Ala Lys Ser Thr Trp Tyr Gly Lys Pro Thr Gly Xaa
    20          25          30
15 Gly Pro Lys Asp Asn Gly Gly Ala Cys Gly Tyr Lys Asp Val Asp Lys
    35          40          45
    Pro Pro Phe Ser Gly Met Thr Gly Cys Gly Asn Thr Pro Ile Phe Lys
      50          55          60
20 Ser Gly Arg Gly Cys Gly Ser Cys Phe Glu Ile Lys Cys Thr Lys Pro
    65          70          75          80
    Glu Ser Cys Ser Gly Glu Pro Val Leu Val His Ile Thr Asp Asp Asn
      85          90          95
25 Glu Glu Pro Ile Ala Ala Tyr His Phe Asp Leu Ser Gly Lys Ala Phe
    100          105          110
30 Gly Ala Met Ala Lys Lys Gly Glu Glu Gln Lys Leu Arg Ser Ala Gly
    115          120          125
    Glu Leu Glu Leu Lys Phe Arg Arg Val Lys Cys Glu Tyr Pro Lys Gly
    130          135          140
35 Thr Lys Val Thr Phe His Val Glu Lys Gly Ser Asn Pro Asn Tyr Leu
    145          150          155          160
    Ala Leu Leu Val Lys Tyr Val Asp Gly Asp Gly Asp Val Val Ala Val
      165          170          175
40 Asp Ile Lys Gln Lys Gly Lys Asp Lys Trp Ile Glu Leu Lys Glu Ser
    180          185          190
45 Trp Gly Ala Val Trp Arg Ile Asp Thr Pro Asp Lys Leu Thr Gly Pro
    195          200          205
    Phe Thr Val Arg Tyr Thr Thr Glu Gly Gly Thr Lys Ala Glu Ala Glu
    210          215          220
50 Asp Val Ile Pro Glu Gly Trp Lys Ala Asp Thr Ala Tyr Glu Ala Lys
    225          230          235          240

```

(2) INFORMATION FOR SEQ ID NO:55:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 723 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 1..720

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

20 ATC GCG AAG GTG CCC CCG GGT CCG AAC ATC ACG GCG ACC TAC GGC GAC
 48
 Ile Ala Lys Val Pro Pro Gly Pro Asn Ile Thr Ala Thr Tyr Gly Asp
 1 5 10 15
 25 AAG TGG CTC GAC GCG AAG AGC ACA TGG TAC GGC AAG CCG ACG GGG GCC
 96
 Lys Trp Leu Asp Ala Lys Ser Thr Trp Tyr Gly Lys Pro Thr Gly Ala
 20 25 30
 30 GGT CCC AAG GAC AAC GGC GGC GCT TGC GGG TAC AAG GAC GTG GAC AAG
 144
 Gly Pro Lys Asp Asn Gly Gly Ala Cys Gly Tyr Lys Asp Val Asp Lys
 35 40 45
 35 CCC CCG TTC AGC GGC ATG ACC GGC TGC GGC AAC ACC CCC ATC TTC AAG
 192
 Pro Pro Phe Ser Gly Met Thr Gly Cys Gly Asn Thr Pro Ile Phe Lys
 50 55 60
 40 TCC GGC CGT GGC TGC GGC TCC TGC TTT GAG ATC AAG TGC ACG AAG CCC
 240
 Ser Gly Arg Gly Cys Gly Ser Cys Phe Glu Ile Lys Cys Thr Lys Pro
 65 70 75 80
 45 GAG GCC TGC TCC GGC GAG CCC GTG GTA GTC CAC ATC ACC GAC GAC AAC
 288
 Glu Ala Cys Ser Gly Glu Pro Val Val Val His Ile Thr Asp Asp Asn
 85 90 95
 50 GAG GAG CCC ATC GCC CCC TAC CAC TTC GAC CTC TCC GGC CAC GCG TTC
 336
 Glu Glu Pro Ile Ala Pro Tyr His Phe Asp Leu Ser Gly His Ala Phe
 100 105 110
 55 GGG GCG ATG GCC AAG AAG GGC GAT GAG CAG AAG CTG CGC ACG GCC GGC
 384
 Gly Ala Met Ala Lys Lys Gly Asp Glu Gln Lys Leu Arg Thr Ala Gly
 115 120 125
 60 GAG CTG GAG CTC CAG TTC CGG CGC GTC AAG TGC AAG TAC CCG GAG GGC
 432
 Glu Leu Glu Leu Gln Phe Arg Arg Val Lys Cys Lys Tyr Pro Glu Gly
 130 135 140

ACC AAG GTG ACC TTC CAC GTG GAG AAG GGG TCC AAC CCC AAC TAC CTG
 480
 Thr Lys Val Thr Phe His Val Glu Lys Gly Ser Asn Pro Asn Tyr Leu
 145 150 155 160
 5 GCG CTG CTT GTG AAG TAC GTT AAC GGC GAC GGA GAC GTG GTG GCG GTG
 528
 Ala Leu Leu Val Lys Tyr Val Asn Gly Asp Gly Asp Val Val Ala Val
 165 170 175
 10 GAC ATC AAG GAG AAG GGC AAG GAC AAG TGG ATC GAG CTC AAG GAG TCG
 576
 Asp Ile Lys Glu Lys Gly Lys Asp Lys Trp Ile Glu Leu Lys Glu Ser
 180 185 190
 15 TGG GGA GCC ATC TGG AGG ATC GAC ACT CCC GAC AAG CTC ACG GGC CCC
 624
 Trp Gly Ala Ile Trp Arg Ile Asp Thr Pro Asp Lys Leu Thr Gly Pro
 195 200 205
 20 TTC ACC GTC CGC TAC ACC ACC GAG GGC GGC ACC AAG ACC GAA GCC GAG
 672
 Phe Thr Val Arg Tyr Thr Thr Glu Gly Gly Thr Lys Thr Glu Ala Glu
 210 215 220
 25 GAC GTC ATC CCT GAG GGC TGG AAG GCC GAC ACC AGC TAC GAG TCC AAG
 720
 Asp Val Ile Pro Glu Gly Trp Lys Ala Asp Thr Ser Tyr Glu Ser Lys
 225 230 235 240
 30 TGA
 723

35 (2) INFORMATION FOR SEQ ID NO:56:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 240 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

45 Ile Ala Lys Val Pro Pro Gly Pro Asn Ile Thr Ala Thr Tyr Gly Asp
 1 5 10 15
 50 Lys Trp Leu Asp Ala Lys Ser Thr Trp Tyr Gly Lys Pro Thr Gly Ala
 20 25 30
 Gly Pro Lys Asp Asn Gly Gly Ala Cys Gly Tyr Lys Asp Val Asp Lys
 35 40 45
 55 Pro Pro Phe Ser Gly Met Thr Gly Cys Gly Asn Thr Pro Ile Phe Lys
 50 55 60
 Ser Gly Arg Gly Cys Gly Ser Cys Phe Glu Ile Lys Cys Thr Lys Pro
 65 70 75 80
 60 Glu Ala Cys Ser Gly Glu Pro Val Val Val His Ile Thr Asp Asp Asn
 85 90 95
 Glu Glu Pro Ile Ala Pro Tyr His Phe Asp Leu Ser Gly His Ala Phe

				100					105								
	Gly	Ala	Met	Ala	Lys	Lys	Gly	Asp	Glu	Gln	Lys	Leu	Arg	Thr	Ala	Gly	
			115					120					125				
5	Glu	Leu	Glu	Leu	Gln	Phe	Arg	Arg	Val	Lys	Cys	Lys	Tyr	Pro	Glu	Gly	
		130					135					140					
	Thr	Lys	Val	Thr	Phe	His	Val	Glu	Lys	Gly	Ser	Asn	Pro	Asn	Tyr	Leu	
10	145					150					155					160	
	Ala	Leu	Leu	Val	Lys	Tyr	Val	Asn	Gly	Asp	Gly	Asp	Val	Val	Ala	Val	
					165					170					175		
15	Asp	Ile	Lys	Glu	Lys	Gly	Lys	Asp	Lys	Trp	Ile	Glu	Leu	Lys	Glu	Ser	
				180					185					190			
	Trp	Gly	Ala	Ile	Trp	Arg	Ile	Asp	Thr	Pro	Asp	Lys	Leu	Thr	Gly	Pro	
			195					200					205				
20	Phe	Thr	Val	Arg	Tyr	Thr	Thr	Glu	Gly	Gly	Thr	Lys	Thr	Glu	Ala	Glu	
		210					215					220					
	Asp	Val	Ile	Pro	Glu	Gly	Trp	Lys	Ala	Asp	Thr	Ser	Tyr	Glu	Ser	Lys	
25	225					230					235					240	

(2) INFORMATION FOR SEQ ID NO:57:

```

30      (i) SEQUENCE CHARACTERISTICS:
          (A) LENGTH: 240 amino acids
          (B) TYPE: amino acid
          (D) TOPOLOGY: linear

35      (ii) MOLECULE TYPE: peptide

          (v) FRAGMENT TYPE: internal

40

      (ix) FEATURE:
          (A) NAME/KEY:
          (B) LOCATION: 45
          (D) OTHER INFORMATION: /note= "Xaa is Asn or Asp"

45      (ix) FEATURE:
          (A) NAME/KEY:
          (B) LOCATION: 144
          (D) OTHER INFORMATION: /note= "Xaa is Asp or Gly"

50      (ix) FEATURE:
          (A) NAME/KEY:
          (B) LOCATION: 154
          (D) OTHER INFORMATION: /note= "Xaa is Gly or Ala"

55      (ix) FEATURE:
          (A) NAME/KEY:
          (B) LOCATION: 187
          (D) OTHER INFORMATION: /note= "Xaa is Ile or Thr"

60      (ix) FEATURE:
          (A) NAME/KEY:
          (B) LOCATION: 223
          (D) OTHER INFORMATION: /note= "Xaa is Val or Phe"

```


(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

5 Ile Ala Lys Val Pro Pro Gly Pro Asn Ile Thr Ala Glu Tyr Gly Asp
 1 5 10 15
 Lys Trp Leu Asp Ala Lys Ser Thr Trp Tyr Gly Lys Pro Thr Gly Ala
 20 25 30
 10 Gly Pro Lys Asp Asn Gly Gly Ala Cys Gly Tyr Lys Xaa Val Asp Lys
 35 40 45
 Ala Pro Phe Asn Gly Met Thr Gly Cys Gly Asn Thr Pro Ile Phe Lys
 50 55 60
 15 Asp Gly Arg Gly Cys Gly Ser Cys Phe Glu Ile Lys Cys Thr Lys Pro
 65 70 75 80
 Glu Ser Cys Ser Gly Glu Ala Val Thr Val Thr Ile Thr Asp Asp Asn
 85 90 95
 Glu Glu Pro Ile Ala Pro Tyr His Phe Asp Leu Ser Gly His Ala Phe
 100 105 110
 25 Gly Ser Met Ala Lys Lys Gly Glu Glu Gln Lys Leu Arg Ser Ala Gly
 115 120 125
 Glu Leu Glu Leu Gln Phe Arg Arg Val Lys Cys Lys Tyr Pro Asp Xaa
 130 135 140
 30 Thr Lys Pro Thr Phe His Val Glu Lys Xaa Ser Asn Pro Asn Tyr Leu
 145 150 155 160
 Ala Ile Leu Val Lys Tyr Val Asp Gly Asp Gly Asp Val Val Ala Val
 165 170 175
 35 Asp Ile Lys Glu Lys Gly Lys Asp Lys Trp Xaa Glu Leu Lys Glu Ser
 180 185 190
 40 Trp Gly Ala Val Trp Arg Ile Asp Thr Pro Asp Lys Leu Thr Gly Pro
 195 200 205
 Phe Thr Val Arg Tyr Thr Thr Glu Gly Gly Thr Lys Ser Glu Xaa Glu
 210 215 220
 45 Asp Val Ile Pro Glu Gly Trp Lys Ala Asp Thr Ser Tyr Ser Ala Lys
 225 230 235 240

50 (2) INFORMATION FOR SEQ ID NO:58:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 240 amino acids
 (B) TYPE: amino acid
 55 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

60 (v) FRAGMENT TYPE: internal

- (ix) FEATURE:
 (A) NAME/KEY:

(B) LOCATION: 199

(D) OTHER INFORMATION: /note= "Xaa is Val or Ile"

(ix) FEATURE:

5

(A) NAME/KEY:

(B) LOCATION: 235

(D) OTHER INFORMATION: /note= "Xaa is Ala or Thr"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

10

Ile	Pro	Lys	Val	Pro	Pro	Gly	Pro	Asn	Ile	Thr	Ala	Thr	Tyr	Gly	Asp
1				5					10					15	

15

Lys	Trp	Leu	Asp	Ala	Lys	Ser	Thr	Trp	Tyr	Gly	Lys	Pro	Thr	Gly	Ala
			20					25					30		

Gly	Pro	Lys	Asp	Asn	Gly	Gly	Ala	Cys	Gly	Tyr	Lys	Asp	Val	Asp	Lys
		35					40					45			

20

Ala	Pro	Phe	Asn	Gly	Met	Thr	Gly	Cys	Gly	Asn	Thr	Pro	Ile	Phe	Lys
	50					55					60				

Asp	Gly	Arg	Gly	Cys	Gly	Ser	Cys	Phe	Glu	Ile	Lys	Cys	Thr	Lys	Pro
65					70					75					80

25

Glu	Ser	Cys	Ser	Gly	Glu	Ala	Val	Thr	Val	His	Ile	Thr	Asp	Asp	Asn
				85					90					95	

30

Glu	Glu	Pro	Ile	Ala	Pro	Tyr	His	Phe	Asp	Leu	Ser	Gly	His	Ala	Phe
			100					105					110		

Gly	Ser	Met	Ala	Lys	Lys	Gly	Glu	Glu	Gln	Lys	Leu	Arg	Ser	Ala	Gly
		115					120					125			

	Glu	Leu	Glu	Leu	Gln	Phe	Arg	Arg	Val	Lys	Cys	Lys	Tyr	Pro	Glu	Gly
	130						135					140				
5	Thr	Lys	Val	Thr	Phe	His	Val	Glu	Lys	Gly	Ser	Asn	Pro	Asn	Tyr	Leu
	145					150					155					160
	Ala	Leu	Leu	Val	Lys	Tyr	Val	Asp	Gly	Asp	Gly	Asp	Val	Val	Ala	Val
					165					170					175	
10	Asp	Ile	Lys	Glu	Lys	Gly	Lys	Asp	Lys	Trp	Ile	Ala	Leu	Lys	Glu	Ser
				180					185					190		
	Trp	Gly	Ala	Ile	Trp	Arg	Xaa	Asp	Thr	Pro	Asp	Lys	Leu	Thr	Gly	Pro
			195					200					205			
15	Phe	Thr	Val	Arg	Tyr	Thr	Thr	Glu	Gly	Gly	Thr	Lys	Ser	Glu	Val	Glu
		210					215					220				
20	Asp	Val	Ile	Pro	Glu	Gly	Trp	Lys	Ala	Asp	Xaa	Ser	Tyr	Glu	Ser	Lys
	225					230					235					240

(2) INFORMATION FOR SEQ ID NO:59:

25	(i) SEQUENCE CHARACTERISTICS:
	(A) LENGTH: 240 amino acids
	(B) TYPE: amino acid
	(D) TOPOLOGY: linear
30	(ii) MOLECULE TYPE: peptide
	(v) FRAGMENT TYPE: internal
35	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:
40	Ile Ala Lys Val Pro Pro Gly Pro Asn Ile Thr Ala Thr Tyr Gly Asp
	1 5 10 15
	Lys Trp Leu Asp Ala Lys Ser Thr Trp Tyr Gly Lys Pro Thr Gly Ala
	20 25 30
45	Gly Pro Lys Asp Asn Gly Gly Ala Cys Gly Tyr Lys Asp Val Asp Lys
	35 40 45
	Pro Pro Phe Ser Gly Met Thr Gly Cys Gly Asn Thr Pro Ile Phe Lys
	50 55 60
50	Ser Gly Arg Gly Cys Gly Ser Cys Phe Glu Ile Lys Cys Thr Lys Pro
	65 70 75 80
	Glu Ala Cys Ser Gly Glu Pro Val Val Val His Ile Thr Asp Asp Asn
	85 90 95
55	Glu Glu Pro Ile Ala Pro Tyr His Phe Asp Leu Ser Gly His Ala Phe
	100 105 110
60	Gly Ala Met Ala Lys Lys Gly Asp Glu Gln Lys Leu Arg Thr Ala Gly
	115 120 125
	Glu Leu Glu Leu Gln Phe Arg Arg Val Lys Cys Lys Tyr Pro Glu Gly
	130 135 140

Thr Lys Val Thr Phe His Val Glu Lys Gly Ser Asn Pro Asn Tyr Leu
 145 150 155 160
 5 Ala Leu Leu Val Lys Tyr Val Asn Gly Asp Gly Asp Val Val Ala Val
 165 170 175
 Asp Ile Lys Glu Lys Gly Lys Asp Lys Trp Ile Glu Leu Lys Glu Ser
 180 185 190
 10 Trp Gly Ala Ile Trp Arg Ile Asp Thr Pro Asp Lys Leu Thr Gly Pro
 195 200 205
 Phe Thr Val Arg Tyr Thr Thr Glu Gly Gly Thr Lys Thr Glu Ala Glu
 210 215 220
 15 Asp Val Ile Pro Glu Gly Trp Lys Ala Asp Thr Ser Tyr Glu Ser Lys
 225 230 235 240

(2) INFORMATION FOR SEQ ID NO:60:

20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 240 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 25 (ii) MOLECULE TYPE: peptide
 (v) FRAGMENT TYPE: internal
 30 (ix) FEATURE:
 (A) NAME/KEY:
 (B) LOCATION: 88
 35 (D) OTHER INFORMATION: /note= "Xaa is Val or Ile"
 (ix) FEATURE:
 (A) NAME/KEY:
 (B) LOCATION: 90
 40 (D) OTHER INFORMATION: /note= "Xaa is Val or Ile"
 (ix) FEATURE:
 (A) NAME/KEY:
 (B) LOCATION: 180
 45 (D) OTHER INFORMATION: /note= "Xaa is Gln or Glu"
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:
 50 Ile Ala Lys Val Pro Pro Gly Pro Asn Ile Thr Ala Thr Tyr Gly Asp
 1 5 10 15
 Lys Trp Leu Asp Ala Lys Ser Thr Trp Tyr Gly Lys Pro Thr Gly Xaa
 20 25 30
 55 Gly Pro Lys Asp Asn Gly Gly Ala Cys Gly Tyr Lys Asp Val Asp Lys
 35 40 45
 Pro Pro Phe Ser Gly Met Thr Gly Cys Gly Asn Thr Pro Ile Phe Lys
 50 55 60
 60 Ser Gly Arg Gly Cys Gly Ser Cys Phe Glu Ile Lys Cys Thr Lys Pro
 65 70 75 80
 Glu Ser Cys Ser Gly Glu Pro Xaa Leu Xaa His Ile Thr Asp Asp Asn

	85	90	95
	Glu Glu Pro Ile Ala Ala Tyr His Phe Asp Leu Ser Gly Lys Ala Phe		
	100	105	110
5	Gly Ala Met Ala Lys Lys Gly Glu Glu Gln Lys Leu Arg Ser Ala Gly		
	115	120	125
	Glu Leu Glu Leu Lys Phe Arg Arg Val Lys Cys Glu Tyr Pro Lys Gly		
10	130	135	140
	Thr Lys Val Thr Phe His Val Glu Lys Gly Ser Asn Pro Asn Tyr Leu		
	145	150	155
15	Ala Leu Leu Val Lys Tyr Val Asp Gly Asp Gly Asp Val Val Ala Val		
	165	170	175
	Asp Ile Lys Xaa Lys Gly Lys Asp Lys Trp Ile Glu Leu Lys Glu Ser		
20	180	185	190
	Trp Gly Ala Val Trp Arg Ile Asp Thr Pro Asp Lys Leu Thr Gly Pro		
	195	200	205
	Phe Thr Val Arg Tyr Thr Thr Glu Gly Gly Thr Lys Ala Glu Ala Glu		
25	210	215	220
	Asp Val Ile Pro Glu Gly Trp Lys Ala Asp Thr Ala Tyr Glu Ala Lys		
	225	230	235
			240

30 (2) INFORMATION FOR SEQ ID NO:61:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

45 Val Glu Lys Gly Ser Asn Pro Asn Tyr Leu Ala Leu Leu Val Lys Tyr
 1 5 10 15

Val Asp Gly Asp
 20

50

(2) INFORMATION FOR SEQ ID NO:62:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

55

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

Val Glu Lys Gly Ser Asn Pro Asn Tyr Leu Ala Leu Leu Val Lys Tyr
 1 5 10 15

5 Val Asn Gly Asp
 20

(2) INFORMATION FOR SEQ ID NO:63:

10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: peptide
 (v) FRAGMENT TYPE: internal

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

Gly Asp Val Val Ala Val Asp Ile Lys Glu Lys Gly Lys Asp Lys Trp
 1 5 10 15

25 Ile Ala Leu Lys
 20

(2) INFORMATION FOR SEQ ID NO:64:

30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: peptide
 (v) FRAGMENT TYPE: internal

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

Gly Asp Val Val Ala Val Asp Ile Lys Gln Lys Gly Lys Asp Lys Trp
 1 5 10 15

45 Ile Glu Leu Lys
 20

50 (2) INFORMATION FOR SEQ ID NO:65:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

55 (ii) MOLECULE TYPE: peptide
 (v) FRAGMENT TYPE: internal

60 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

Glu Ser Trp Gly Ala Ile Trp Arg Ile Asp Thr Pro Asp Lys Leu Thr
 1 5 10 15

Gly Pro Phe Thr
 20

5

(2) INFORMATION FOR SEQ ID NO:66:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: peptide

15

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

Glu Ser Trp Gly Ala Ile Trp Arg Val Asp Thr Pro Asp Lys Leu Thr
 1 5 10 15

Gly Pro Phe Thr
 20

25

(2) INFORMATION FOR SEQ ID NO:67:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: peptide

35

(v) FRAGMENT TYPE: internal

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

Thr Glu Ala Glu Asp Val Ile Pro Glu Gly Trp Lys Ala Asp Thr Ser
 1 5 10 15

Tyr Glu Ser Lys
 20

45

(2) INFORMATION FOR SEQ ID NO:68:

50

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

55

(ii) MOLECULE TYPE: peptide

(v) FRAGMENT TYPE: internal

60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

Ala Glu Ala Glu Asp Val Ile Pro Glu Gly Trp Lys Ala Asp Thr Ala

1 5 10 15

Tyr Glu Ala Lys
20

5

(2) INFORMATION FOR SEQ ID NO:69:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: peptide

15

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

20

Ser Glu Val Glu Asp Val Ile Pro Glu Gly Trp Lys Ala Asp Ala Ser
1 5 10 15

Tyr Glu Ser Lys
20

25

(2) INFORMATION FOR SEQ ID NO:70:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

30

(ii) MOLECULE TYPE: peptide

35

(v) FRAGMENT TYPE: internal

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

40

Ser Glu Val Glu Asp Val Ile Pro Glu Gly Trp Lys Ala Asp Thr Ser
1 5 10 15

Tyr Glu Ser Lys
20

45

(2) INFORMATION FOR SEQ ID NO:71:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

50

55

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

60

GGGTCTAGAG GTACCGTCCG ATCGATCATT
30

(2) INFORMATION FOR SEQ ID NO:72:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 13 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

AATGATCGAT GCT
13

(2) INFORMATION FOR SEQ ID NO:73:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:

GGGTCTAGAG GTACCGTCCG
20

(2) INFORMATION FOR SEQ ID NO:74:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:74:

CCCTGCAGAT TATTTGAGAT CTTGAG
26

(2) INFORMATION FOR SEQ ID NO:75:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 29 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:75:

CCCTGCAGTC ATGCTCACTT GGCCGAGTA
29

5 (2) INFORMATION FOR SEQ ID NO:76:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 base pairs
 (B) TYPE: nucleic acid
10 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:76:

20 GAGTACGGCG ACAAGTGGC
19

(2) INFORMATION FOR SEQ ID NO:77:

- (i) SEQUENCE CHARACTERISTICS:
25 (A) LENGTH: 18 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:77:

35 TTCGAGATCA AGTGCACC
18

(2) INFORMATION FOR SEQ ID NO:78:

- 40 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 16 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
45 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:78:

GTGACAGCCT CGCCGG
16

5

(2) INFORMATION FOR SEQ ID NO:79:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: cDNA

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:79:

GGGAATTCCA TGGCGAAGAA GGGC
24

20

(2) INFORMATION FOR SEQ ID NO:80:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 16 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25

(ii) MOLECULE TYPE: cDNA

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:80:

GTGCCGTCCG GGTACT
16

35

(2) INFORMATION FOR SEQ ID NO:81:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 16 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: cDNA

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:81:

CCGTCGACGT ACTTCA
16

55

(2) INFORMATION FOR SEQ ID NO:82:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:82:

GGAGTCGTGG GGAGCAGTC
19

(2) INFORMATION FOR SEQ ID NO:83:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 19 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:83:

GGGTCTAGAG GTACCGTCC
19

(2) INFORMATION FOR SEQ ID NO:84:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:84:

TTGGATCCTA CGGCAAGCCG ACCGGC
26

(2) INFORMATION FOR SEQ ID NO:85:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 28 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:85:

TTGGATCCAT CCCGAAGGTG CCCCCGGG
28

5 (2) INFORMATION FOR SEQ ID NO:86:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 base pairs
 (B) TYPE: nucleic acid
10 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:86:

20 AGGTGACCTT CCACGTCG
18

(2) INFORMATION FOR SEQ ID NO:87:

- (i) SEQUENCE CHARACTERISTICS:
25 (A) LENGTH: 27 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:87:

35 TTGGATCCTG GCGCTGCTGG TGAAGTA
27

(2) INFORMATION FOR SEQ ID NO:88:

- 40 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 26 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
45 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:88:
TTGAATTCAT CCCGAAGGTG CCCCCG
26

5 (2) INFORMATION FOR SEQ ID NO:89:

10 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:89:
TTGGTACCTC ACTTGGACTC GTAGCT
26

20 (2) INFORMATION FOR SEQ ID NO:90:

25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 25 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: cDNA

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:90:
CCGAATTCGT GGAGAAGGGG TCCAA
25

40 (2) INFORMATION FOR SEQ ID NO:91:

45 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

50 (ii) MOLECULE TYPE: cDNA

(ix) FEATURE:
(A) NAME/KEY: Modified-site
(B) LOCATION: 22
55 (D) OTHER INFORMATION: /note= "Xaa is Iosine"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:91:

TTAGGATCCT CACTTATCAT ANGACGTATC
30

5

(2) INFORMATION FOR SEQ ID NO:92:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: cDNA

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:92:

TTGAATTCCT TGTCATTGCC CTTCTG
26

20

(2) INFORMATION FOR SEQ ID NO:93:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

25

30

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:93:

AAGAATTCCT TCTGCTTGAT GTCCAC
26

35

(2) INFORMATION FOR SEQ ID NO:94:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40

45

(ii) MOLECULE TYPE: cDNA

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:94:

ATGAATTCGA GTCGTGGGGA GCCGTC
26

55

(2) INFORMATION FOR SEQ ID NO:95:

(i) SEQUENCE CHARACTERISTICS:

5 (A) LENGTH: 26 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:95:

15 ATGAATTCGT CTGGAGGATC GACACC

26

(2) INFORMATION FOR SEQ ID NO:96:

20 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

25 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:96:

ATGAATTCAT CGCAAAGGTT CCCCCC

26

35 (2) INFORMATION FOR SEQ ID NO:97:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27 base pairs

40 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:97:

TTTGGATCCT CACTTGGACT CGTAGCT

50 27

(2) INFORMATION FOR SEQ ID NO:98:

55 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

60 (ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:98:

TTGAATTCTC GCGAAGGTGC CCCCC
25

Claims

1. An isolated peptide of *Lol p I* or an isolated portion thereof, said peptide
or portion thereof comprising at least one T cell epitope of *Lol p I*, said peptide
5 comprising an amino acid sequence selected from the group consisting of: LPI-1
(SEQ ID NO: 4).1, LPI-2 (SEQ ID NO: 5), LPI-3 (SEQ ID NO: 6), LPI-4 (SEQ
ID NO: 7), LPI-4.1 (SEQ ID NO: 8), LPI-8 (SEQ ID NO: 12), LPI-10 (SEQ ID
NO: 14), LPI-11 (SEQ ID NO: 15), LPI-13 (SEQ ID NO: 19), LPI-15 (SEQ ID
NO: 21), LPI-16 (SEQ ID NO: 22), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ
10 ID NO: 25), LPI-19 (SEQ ID NO: 26), LPI-22 (SEQ ID NO: 29) and LPI-23
(SEQ ID NO: 30), all as shown in Fig. 2.
2. An isolated peptide of *Lol p I* or an isolated portion thereof, said peptide
or portion thereof comprising at least one T cell epitope of *Lol p I*, said peptide
15 having an amino acid sequence selected from the group consisting of: LPI-16.2
(SEQ ID NO: 31), LPI-16.3 (SEQ ID NO: 32), LPI-16.4 (SEQ ID NO: 33), LPI-
16.5 (SEQ ID NO: 34), LPI-16.6 (SEQ ID NO: 35), LPI-16.7 (SEQ ID NO: 36),
LPI-16.9 (SEQ ID NO: 37), LPI-16.10 (SEQ ID NO: 38), LPI-18.5 (SEQ ID
NO: 39), LPI-18.6 (SEQ ID NO: 40), LPI-18.7 (SEQ ID NO: 41), LPI-18.8
20 (SEQ ID NO: 42), LPI-20.2 (SEQ ID NO: 43), LPI-20.3 (SEQ ID NO: 44), LPI-
20.4 (SEQ ID NO: 45), LPI-20.5 (SEQ ID NO: 46), LPI-20.6 (SEQ ID NO: 47),
LPI-23.1 (SEQ ID NO: 48), LPI-23.2 (SEQ ID NO: 49), and LPI-23.4 (SEQ ID
NO: 50), all as shown in Fig. 4.
- 25 3. An isolated peptide or portion thereof according to claim 1, wherein said
portion of a peptide has a mean T cell stimulation index approximately equivalent
to or greater than the mean T cell stimulation index of the corresponding peptide
shown in Fig. 3.
- 30 4. An isolated peptide or portion thereof of claim 1 or 2 which comprises at
least two T cell epitopes.
5. An isolated peptide or portion thereof of claim 1 or 2 which induces T cell
nonresponsiveness or modifies the lymphokine secretion profile of appropriate T
35 cell subpopulations.

6. An isolated peptide or portion thereof of claim 1 or 2 which, when administered to an individual sensitive to an allergen of the family, Poacea induces T cell anergy or modifies the lymphokine secretion profile of appropriate T cell populations.
- 5
7. A portion of an isolated peptide of claim 1 or 2 which has a mean T cell stimulation index of at least 3.5.
8. An isolated peptide or a portion thereof of claim 1 or 2 which does not
10 bind immunoglobulin E specific for *Lol p I* in a substantial percentage of individuals sensitive to *Lol p I*, or if binding of the peptide or portion thereof to said immunoglobulin E occurs, such binding does not result in release of mediators from mast cells or basophils in a substantial percentage of individuals sensitive to *Lol p I*.
- 15
9. An isolated peptide of claim 1 or 2 which binds immunoglobulin E to a substantially lesser extent than purified native *Lol p I* binds immunoglobulin E.
10. An isolated peptide or portion thereof of claim 1 or 2 which, when
20 administered to an individual sensitive to *Lol p I* allergen, modifies the allergic response of the individual to ryegrass pollen allergen.
11. An isolated peptide or portion thereof of claim 1 or 2 which, when
25 administered to an individual sensitive to an allergen of the family Poacea, modifies the allergic response of the individual to said allergen.
12. A portion of an isolated peptide of claim 1 or 2 wherein said portion comprises at least 15 amino acid residues.
- 30
13. An isolated nucleic acid having a sequence encoding all or a portion of a peptide of claim 1 or 2.
14. A functional equivalent of a nucleic acid sequence encoding all or a
35 portion of a peptide of claim 1 or 2.

15. An isolated peptide that is immunologically cross-reactive with T cells reactive with a peptide of claim 1 or 2.
16. An isolated peptide or portion thereof of *Lol p I*, said peptide or portion thereof comprising at least one T cell epitope of *Lol p I*, said peptide having a positivity index of at least about 100 and mean T cell stimulation index of at least about 3.0 determined in a population of individuals sensitive to said protein allergen.
17. An isolated peptide or portion thereof of claim 16 wherein said population of individuals is at least thirty individuals.
18. An isolated peptide or portion thereof of claim 17 wherein said population of individuals is at least thirty-five individuals.
19. An isolated peptide or portion thereof of claim 17 wherein said mean T cell stimulation index is at least about 4.0.
20. An isolated peptide or portion thereof of claim 17 wherein said mean T cell stimulation index is at least about 6.0.
21. A peptide or portion thereof of claim 17 wherein said peptide is selected from the group consisting of: LPI-2 (SEQ ID NO: 5), LPI-11 (SEQ ID NO: 15), LPI-13 (SEQ ID NO: 19), LPI-15 (SEQ ID NO: 21), LPI-16 (SEQ ID NO: 22), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), LPI-22 (SEQ ID NO: 29) and LPI-23 (SEQ ID NO: 30).
22. An isolated peptide of *Lol p I*, or a portion thereof wherein said peptide is selected from the group consisting of: LPI-1.1 (SEQ ID NO: 4), LPI-2 (SEQ ID NO: 5), LPI-3 (SEQ ID NO: 6), LPI-4 (SEQ ID NO: 7), LPI-4.1 (SEQ ID NO: 8), LPI-8 (SEQ ID NO: 12), LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-13 (SEQ ID NO: 19), LPI-15 (SEQ ID NO: 21), LPI-16 (SEQ ID NO: 22), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), LPI-19 (SEQ ID NO: 26), LPI-22 (SEQ ID NO: 29), LPI-23 (SEQ ID NO: 30), LPI-18.5 (SEQ ID NO: 39), LPI-18.6 (SEQ ID NO: 40), LPI-18.7 (SEQ ID NO: 41), LPI-

18.8 (SEQ ID NO: 42), LPI-20.2 (SEQ ID NO: 43), LPI-20.3 (SEQ ID NO: 44), LPI-20.4 (SEQ ID NO: 45), LPI-20.5 (SEQ ID NO: 46), LPI-20.6 (SEQ ID NO: 47), LPI-23.1 (SEQ ID NO: 48), LPI-23.2 (SEQ ID NO: 49), and LPI-23.4 (SEQ ID NO: 50) or portion thereof.

5

23. A modified peptide or a modified portion of a peptide of claim 22.

24. A modified peptide of claim 23 wherein said peptide is selected from the group consisting of: LPI-16.2 (SEQ ID NO: 31), LPI-16.3 (SEQ ID NO: 32),
10 LPI-16.4 (SEQ ID NO: 33), LPI-16.5 (SEQ ID NO: 34), LPI-16.6 (SEQ ID NO: 35), LPI-16.7 (SEQ ID NO: 36), LPI-16.9 (SEQ ID NO: 37), and LPI-16.10 (SEQ ID NO: 38), all as shown in Fig. 4.

25. A modified peptide or a modified portion of a peptide of claim 23 or 24
15 which does not bind immunoglobulin E specific for *Lol p I* in a substantial percentage of individuals sensitive to *Lol p I*, or if binding of the peptide or portion thereof to said immunoglobulin E occurs, such binding does not result in release of mediators from mast cells or basophils in a substantial percentage of individuals sensitive to *Lol p I*.

20

26. A modified peptide or a modified portion of a peptide of claim 23 or 24 which modifies, in an individual sensitive to *Lol p I* or an immunologically related allergen, the allergic response of the individual to *Lol p I* allergen or said related allergen.

25

27. An isolated peptide comprising at least two regions, each region comprising at least one T cell epitope of *Lol p I*, said regions each comprising all or a portion of an amino acid sequence selected from the group consisting of:
LPI-1.1 (SEQ ID NO: 4), LPI-2 (SEQ ID NO: 5), LPI-3 (SEQ ID NO: 6), LPI-4
30 (SEQ ID NO: 7), LPI-4.1 (SEQ ID NO: 8), LPI-8 (SEQ ID NO: 12), LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-13 (SEQ ID NO: 19), LPI-15 (SEQ ID NO: 21), LPI-16 (SEQ ID NO: 22), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), LPI-23 (SEQ ID NO: 30), LPI-16.2 (SEQ ID NO: 31), LPI-16.3 (SEQ ID NO: 32),
35 LPI-16.4 (SEQ ID NO: 33), LPI-16.5 (SEQ ID NO: 34), LPI-16.6 (SEQ ID

NO: 35), LPI-16.7 (SEQ ID NO: 36), LPI-16.9 (SEQ ID NO: 37), LPI-16.10 (SEQ ID NO: 38), LPI-18.5 (SEQ ID NO: 39), LPI-18.6 (SEQ ID NO: 40), LPI-18.7 (SEQ ID NO: 41), LPI-18.8 (SEQ ID NO: 42), LPI-20.2 (SEQ ID NO: 43), LPI-20.3 (SEQ ID NO: 44), LPI-20.4 (SEQ ID NO: 45), LPI-20.5 (SEQ ID NO: 46), LPI-20.6 (SEQ ID NO: 47), LPI-23.1 (SEQ ID NO: 48), LPI-23.2 (SEQ ID NO: 49), and LPI-23.4 (SEQ ID NO: 50).

28. An isolated peptide of claim 27 wherein said regions comprise an amino acid sequence selected from the group consisting of: LPI-3 (SEQ ID NO: 6), LPI-4.1 (SEQ ID NO: 8), LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), LPI-23 (SEQ ID NO: 30), LPI-16.2 (SEQ ID NO: 31), LPI-16.3 (SEQ ID NO: 32), LPI-16.4 (SEQ ID NO: 33), LPI-16.5 (SEQ ID NO: 34), LPI-16.6 (SEQ ID NO: 35), LPI-16.7 (SEQ ID NO: 36), LPI-16.9 (SEQ ID NO: 37), LPI-16.10 (SEQ ID NO: 38), LPI-18.5 (SEQ ID NO: 39), LPI-18.6 (SEQ ID NO: 40), LPI-18.7 (SEQ ID NO: 41), LPI-18.8 (SEQ ID NO: 42), LPI-20.2 (SEQ ID NO: 43), LPI-20.3 (SEQ ID NO: 44), LPI-20.4 (SEQ ID NO: 45), LPI-20.5 (SEQ ID NO: 46), LPI-20.6 (SEQ ID NO: 47), LPI-23.1 (SEQ ID NO: 48), LPI-23.2 (SEQ ID NO: 49), and LPI-23.4 (SEQ ID NO: 50), or a portion thereof containing at least two *Lol p I* epitopes.

29. An isolated peptide of *Lol p I*, wherein said peptide comprises a combination of regions selected from the group consisting of:

LPI-3 (SEQ ID NO: 6), LPI-4.1 (SEQ ID NO: 8), LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), LPI-16 (SEQ ID NO: 22), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), and LPI-23 (SEQ ID NO: 30);

LPI-3 (SEQ ID NO: 6), LPI-4.1 (SEQ ID NO: 8), LPI-10 (SEQ ID NO: 14), and LPI-11 (SEQ ID NO: 15);

LPI-3 (SEQ ID NO: 6), LPI-4.1 (SEQ ID NO: 8), LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), and LPI-16 (SEQ ID NO: 22);

LPI-3 (SEQ ID NO: 6), LPI-4.1 (SEQ ID NO: 8), LPI-10 (SEQ ID NO: 14),
LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), and LPI-16.1 (SEQ ID
NO: 23);

5 LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO:
21), and LPI-16.1 (SEQ ID NO: 23);

LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO:
21), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), and LPI-20 (SEQ
ID NO: 27);

10 LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO:
21), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID
NO: 27), LPI-22 (SEQ ID NO: 29) and LPI-23 (SEQ ID NO: 30);

LPI-15 (SEQ ID NO: 21), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO:
25), and LPI-20 (SEQ ID NO: 27);

15 LPI-15 (SEQ ID NO: 21), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO:
25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), and LPI-23 (SEQ
ID NO: 30);

LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO:
29), and LPI-23 (SEQ ID NO: 30);

LPI-18 (SEQ ID NO: 25) and LPI-20 (SEQ ID NO: 27);

20 LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27) and LPI-23 (SEQ ID
NO: 30);

LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27) and LPI-16.1 (SEQ ID
NO: 23);

25 LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO:
30) and LPI-16.1 (SEQ ID NO: 23);

LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO:
30), LPI-16.1 (SEQ ID NO: 23) and LPI-11 (SEQ ID NO: 15);

LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO:
30), LPI-16.1 (SEQ ID NO: 23) and LPI-4.1 (SEQ ID NO: 8);

30 LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO:
30), LPI-16 (SEQ ID NO: 23).1, LPI-4.1 (SEQ ID NO: 8) and LPI-22 (SEQ
ID NO: 29);

LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO:
30), LPI-16.1 (SEQ ID NO: 23), LPI-11 (SEQ ID NO: 15) and LPI-4.1 (SEQ
ID NO: 8);

35

LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO: 30), LPI-16.1 (SEQ ID NO: 23), LPI-11 (SEQ ID NO: 15), LPI-4.1 (SEQ ID NO: 8) and LPI-22 (SEQ ID NO: 29);

5 LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), and LPI-23 (SEQ ID NO: 30);

LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-16.1 (SEQ ID NO: 23), LPI-22 (SEQ ID NO: 29) and LPI-23 (SEQ ID NO: 30); and

LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-16.1 (SEQ ID NO: 23) and LPI-22 (SEQ ID NO: 29).

10

30. An isolated peptide of *Lol p I*, wherein said peptide comprises a combination of regions selected from the group consisting of:

LPI-16.2 (SEQ ID NO: 31), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);

15 LPI-16.3 (SEQ ID NO: 32), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);

LPI-16.4 (SEQ ID NO: 33), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);

20 LPI-16.5 (SEQ ID NO: 34), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);

LPI-16.6 (SEQ ID NO: 35), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);

LPI-16.7 (SEQ ID NO: 36), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);

25 LPI-16.9 (SEQ ID NO: 37), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30); and

LPI-16.10 (SEQ ID NO: 38), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30).

30 31. A monoclonal antibody, polyclonal antibody, or immunoreactive fragment thereof specifically reactive with a peptide of claim 1 or 2.

32. An isolated peptide produced in a host cell transformed with the nucleic acid of claim 13.

35

33. An isolated peptide produced in a host cell transformed with the nucleic acid of claim 14.
34. An isolated nucleic acid having a sequence encoding a peptide of claim 27
5 or 29.
35. The functional equivalent of an isolated nucleic acid sequence encoding a peptide of claim 27 or 29.
- 10 36. An isolated peptide produced in a host cell transformed with the nucleic acid of claim 34.
37. An expression vector comprising a nucleic acid sequence coding for a peptide of claim 1 or 2.
15
38. An expression vector comprising the functional equivalent of a sequence coding for a peptide of claim 1 or 2.
39. An expression vector comprising a nucleic acid sequence coding for a
20 peptide of claim 27 or 29.
40. An expression vector comprising the functional equivalent of a nucleic acid sequence coding for a peptide of claim 27 or 29.
- 25 41. All or a portion of an isolated peptide of *Lol p I*, said peptide or portion thereof comprising at least one T cell epitope of said protein allergen, said peptide having the formula X_n-Y-Z_m , wherein Y is an amino acid sequence selected from the group consisting of: LPI-1 (SEQ ID NO: 3), LPI-1.1 (SEQ ID NO: 4), LPI-2 (SEQ ID NO: 5), LPI-3 (SEQ ID NO: 6), LPI-4 (SEQ ID NO: 7), LPI-4.1 (SEQ
30 ID NO: 8), LPI-5 (SEQ ID NO: 9), LPI-6 (SEQ ID NO: 10), LPI-7 (SEQ ID NO: 11), LPI-8 (SEQ ID NO: 12), LPI-9 (SEQ ID NO: 13), LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-12 (SEQ ID NO: 17), LPI-13 (SEQ ID NO: 19), LPI-14 (SEQ ID NO: 20), LPI-15 (SEQ ID NO: 21), LPI-16 (SEQ ID NO: 22), LPI-16.1 (SEQ ID NO: 23), LPI-17 (SEQ ID NO: 24), LPI-18 (SEQ
35 ID NO: 25), LPI-19 (SEQ ID NO: 26), LPI-21 (SEQ ID NO: 28), LPI-22 (SEQ

- ID NO: 29), LPI-23 (SEQ ID NO: 30), LPI-16.2 (SEQ ID NO: 31), LPI-16.3 (SEQ ID NO: 32), LPI-16.4 (SEQ ID NO: 33), LPI-16.5 (SEQ ID NO: 34), LPI-16.6 (SEQ ID NO: 35), LPI-16.7 (SEQ ID NO: 36), LPI-16.9 (SEQ ID NO: 37), LPI-16.10 (SEQ ID NO: 38), LPI-18.5 (SEQ ID NO: 39), LPI-18.6 (SEQ ID NO: 40), LPI-18.7 (SEQ ID NO: 41), LPI-18.8 (SEQ ID NO: 42), LPI-20.2 (SEQ ID NO: 43), LPI-20.3 (SEQ ID NO: 44), LPI-20.4 (SEQ ID NO: 45), LPI-20.5 (SEQ ID NO: 46), LPI-20.6 (SEQ ID NO: 47), LPI-23.1 (SEQ ID NO: 48), LPI-23.2 (SEQ ID NO: 49), and LPI-23.4 (SEQ ID NO: 50) wherein X_n are amino acid residues contiguous to the amino terminus of Y in the amino acid sequence of said protein allergen, wherein Z_m are amino acid residues contiguous to the carboxy terminus of Y in the amino acid sequence of said protein allergen, wherein n is 0-30 and wherein m is 0-30.
42. A portion of an isolated peptide of claim 40 wherein the portion comprises at least fifteen amino acid residues.
43. A composition comprising at least one isolated peptide or a portion thereof of claim 1 or 2 and a pharmaceutically acceptable carrier or diluent.
44. A composition comprising at least one isolated peptide or portion thereof of claim 23 or 24 and a pharmaceutically acceptable carrier or diluent.
45. A composition comprising an isolated peptide or portion thereof of claim 27 or 29 and a pharmaceutically acceptable carrier or diluent.
46. Use of a composition of claim 43 in the manufacture of a medicament for treating sensitivity to *Lol p I* protein allergen or an allergen which is immunologically cross-reactive with *Lol p I* protein allergen.
47. Use of a composition of claim 44 in the manufacture of a medicament for treating sensitivity to *Lol p I* protein allergen or an allergen which is immunologically cross-reactive with *Lol p I* protein allergen.

48. Use of at least two compositions of claim 43 in the manufacture of a medicament for treating sensitivity to *Lol p I* protein allergen or an allergen which is immunologically cross-reactive with *Lol p I* protein allergen.

5 49. The use of the composition of claim 46 wherein said immunologically cross-reactive allergen is *Dac g I*, *Poa p I* or *Phl p I*.

10 50. A method of detecting sensitivity to *Lol p I* protein allergen or an immunologically cross-reactive allergen in an individual, comprising combining a blood sample obtained from the individual with at least one peptide of claim 1 or 2, *in vitro*, under conditions appropriate for binding of blood components with the peptide, and determining the extent to which such binding occurs as indicative of sensitivity in the individual to ryegrass pollen allergen or said immunologically cross-reactive allergen.

15 51. A method of claim 50 wherein the extent to which binding occurs is determined by assessing T cell function, T cell proliferation or a combination thereof.

20 52. A composition comprising a pharmaceutically acceptable carrier or diluent and at least two peptides, selected from the group consisting of: LPI-1.1 (SEQ ID NO: 4), LPI-2 (SEQ ID NO: 5), LPI-3 (SEQ ID NO: 6), LPI-4 (SEQ ID NO: 7), LPI-4.1 (SEQ ID NO: 8), LPI-8 (SEQ ID NO: 12), LPI-11 (SEQ ID NO: 15), LPI-13 (SEQ ID NO: 19), LPI-15 (SEQ ID NO: 21), LPI-16 (SEQ ID NO: 22),
25 LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), LPI-23 (SEQ ID NO: 30), LPI-16.2 (SEQ ID NO: 31), LPI-16.3 (SEQ ID NO: 32), LPI-16.4 (SEQ ID NO: 33), LPI-16.5 (SEQ ID NO: 34), LPI-16.6 (SEQ ID NO: 35), LPI-16.7 (SEQ ID NO: 36), LPI-16.9 (SEQ ID NO: 37), LPI-16.10 (SEQ ID NO: 38), LPI-18.5 (SEQ ID NO:
30 39), LPI-18.6 (SEQ ID NO: 40), LPI-18.7 (SEQ ID NO: 41), LPI-18.8 (SEQ ID NO: 42), LPI-20.2 (SEQ ID NO: 43), LPI-20.3 (SEQ ID NO: 44), LPI-20.4 (SEQ ID NO: 45), LPI-20.5 (SEQ ID NO: 46), LPI-20.6 (SEQ ID NO: 47), LPI-23.1 (SEQ ID NO: 48), LPI-23.2 (SEQ ID NO: 49), and LPI-23.4 (SEQ ID NO: 50) and wherein said composition comprises a sufficient percentage of the T cell
35 epitopes of said protein allergen such that T cells of an individual sensitive to *Lol*

p I protein pollen or an immunologically cross-reactive allergen, are tolerized to said at least one protein allergen.

53. A composition of claim 43 comprising a combination of peptides selected
- 5 from the group consisting of:
- LPI-3 (SEQ ID NO: 6), LPI-4.1 (SEQ ID NO: 8), LPI-10 (SEQ ID NO: 14),
LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), LPI-16 (SEQ ID NO:
22), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID
NO: 29), and LPI-23 (SEQ ID NO: 30);
- 10 LPI-3 (SEQ ID NO: 6), LPI-4.1 (SEQ ID NO: 8), LPI-10 (SEQ ID NO: 14),
and LPI-11 (SEQ ID NO: 15);
- LPI-3 (SEQ ID NO: 6), LPI-4.1 (SEQ ID NO: 8), LPI-10 (SEQ ID NO: 14),
LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), and LPI-16 (SEQ ID
NO: 22);
- 15 LPI-3 (SEQ ID NO: 6), LPI-4.1 (SEQ ID NO: 8), LPI-10 (SEQ ID NO: 14),
LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO: 21), and LPI-16.1 (SEQ ID
NO: 23);
- LPI-10 (SEQ ID NO: 14), LPI-11 (SEQ ID NO: 15), LPI-15 (SEQ ID NO:
21), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID
NO: 27), LPI-22 (SEQ ID NO: 29) and LPI-23 (SEQ ID NO: 30);
- 20 LPI-15 (SEQ ID NO: 21), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO:
25), and LPI-20 (SEQ ID NO: 27);
- LPI-15 (SEQ ID NO: 21), LPI-16.1 (SEQ ID NO: 23), LPI-18 (SEQ ID NO:
25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), and LPI-23 (SEQ
ID NO: 30);
- 25 LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO:
29), and LPI-23 (SEQ ID NO: 30);
- LPI-18 (SEQ ID NO: 25) and LPI-20 (SEQ ID NO: 27);
- LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27) and LPI-23 (SEQ ID
NO: 30);
- 30 LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27) and LPI-16.1 (SEQ ID
NO: 23);
- LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO:
30) and LPI-16.1 (SEQ ID NO: 23);

LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO: 30), LPI-16.1 (SEQ ID NO: 23) and LPI-11 (SEQ ID NO: 15);
LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO: 30), LPI-16.1 (SEQ ID NO: 23) and LPI-4.1 (SEQ ID NO: 8);
5 LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO: 30), LPI-16.1 (SEQ ID NO: 23), LPI-4.1 (SEQ ID NO: 8) and LPI-22 (SEQ ID NO: 29);
LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO: 30), LPI-16.1 (SEQ ID NO: 23), LPI-11 (SEQ ID NO: 15) and LPI-4.1 (SEQ ID NO: 8);
10 LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-23 (SEQ ID NO: 30), LPI-16.1 (SEQ ID NO: 23), LPI-11 (SEQ ID NO: 15), LPI-4.1 (SEQ ID NO: 8) and LPI-22 (SEQ ID NO: 29);
LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-22 (SEQ ID NO: 29), and LPI-23 (SEQ ID NO: 30);
15 LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-16.1 (SEQ ID NO: 23), LPI-22 (SEQ ID NO: 29) and LPI-23 (SEQ ID NO: 30); and
LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), LPI-16.1 (SEQ ID NO: 23) and LPI-22 (SEQ ID NO: 29).

20

54. A composition of claim 43 comprising a combination of peptides selected from the group consisting of:

LPI-16.2 (SEQ ID NO: 31), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);
25 LPI-16.3 (SEQ ID NO: 32), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);
LPI-16.4 (SEQ ID NO: 33), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);
LPI-16.5 (SEQ ID NO: 34), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);
30 LPI-16.6 (SEQ ID NO: 35), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);
LPI-16.7 (SEQ ID NO: 36), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30);

LPI-16.9 (SEQ ID NO: 37), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30); and
LPI-16.10 (SEQ ID NO: 38), LPI-18 (SEQ ID NO: 25), LPI-20 (SEQ ID NO: 27), and LPI-23 (SEQ ID NO: 30).

5

55. Use of composition of claim 52, 53 or 54 in the manufacture of a medicament for use in treating sensitivity to *Lol p I* allergen or an immunologically cross-reactive allergen.

- 10 56. A method of designing antigenic fragments of *Lol p I*, which when administered to ryegrass pollen sensitive individuals in sufficient quantity will modify the individual's allergic reaction to ryegrass pollen comprising the steps of:
- (a) recombinantly or synthetically producing peptides of *Lol p I*;
 - (b) examining said peptides for their ability to influence B cell and/or
 - 15 T cell responses in ryegrass pollen sensitive individuals;
 - (c) selecting appropriate peptides which contain epitopes recognized by the cells, and
 - (d) combining epitope-containing regions to include multiple epitopes in one peptide.

20

57. A method of designing antigenic fragments of *Lol p I*, which when administered to ryegrass pollen sensitive individuals in sufficient quantity will modify the individual's allergic reaction to ryegrass pollen comprising the steps of:
- (a) recombinantly or synthetically producing peptides of *Lol p I*;
 - 25 (b) examining said peptides for their ability to influence B cell and/or T cell responses in ryegrass pollen sensitive individuals; and
 - (c) selecting appropriate peptides which contain epitopes recognized by the cells.

30

58. A T cell capable of recognizing a peptide of claim 1 or 2.

59. A receptor of a T cell capable of recognizing a peptide of claim 1 or 2.

35

60. An isolated nucleic acid having a nucleotide sequence coding for *Dac g I*, or the functional equivalent of said nucleotide sequence.

61. An isolated nucleic acid sequence of claim 60 wherein said nucleotide sequence comprises the nucleotide sequence of Fig. 5.
- 5 62. An expression vector comprising a nucleotide sequence coding for *Dac g I*, or the functional equivalent of said nucleotide sequence.
63. A host cell transformed to express a protein encoded by the nucleic acid of claim 60.
- 10 64. Isolated *Dac g I* protein produced in a host cell transformed with the nucleic acid of claim 60.
65. An isolated nucleic acid having a nucleotide sequence coding for *Poa p I*, or
15 the functional equivalent of said nucleotide sequence.
66. An isolated nucleic acid sequence of claim 65 wherein said nucleotide sequence comprises the nucleotide sequence of Fig. 6.
- 20 67. An expression vector comprising a nucleotide sequence coding for *Poa p I*, or the functional equivalent of said nucleotide sequence.
68. A host cell transformed to express a protein encoded by the nucleic acid of claim 65.
- 25 69. Isolated *Poa p I* protein produced in a host cell transformed with the nucleic acid of claim 60.
70. An isolated protein allergen that is immunologically related to *Lol p I*.
- 30 71. An isolated protein allergen of claim 70 wherein said protein allergen is *Dac g I* or *Poa p I*.

```

CAAATTCAAG ACAAG ATG GCG TCC TCC TCG TCG GTG CTC CTG GTG GTG GCG GCG 51
Met Ala Ser Ser Ser Ser Val Leu Leu Val Val Ala
-23 -20 -15

CTG TTC GCC GTG TTC CTG GGC AGC GCG CAT GGC ATC GCG AAG GTA CCA 99
Leu Phe Ala Val Phe Leu Gly Ser Ala His Gly Ile Ala Lys Val Pro
-10 -5 1 5

CCG GGC CCC AAC ATC ACG GCC GAG TAC GGC GAC AAG TGG CTG GAC GCG 147
Pro Gly Pro Asn Ile Thr Ala Glu Tyr Gly Asp Lys Trp Leu Asp Ala
10 15 20

AAG AGC ACC TGG TAT GGC AAG CCG ACC GGC GCC GGT CCC AAG GAC AAC 195
Lys Ser Thr Trp Tyr Gly Lys Pro Thr Gly Ala Gly Pro Lys Asp Asn
25 30 35

GGC GGC GCG TGC GGG TAC AAG GAC GTT GAC AAG GCG CCG TTC AAC GCG 243
Gly Gly Ala Cys Gly Tyr Lys Asp Val Asp Lys Ala Pro Phe Asn Gly
40 45 50

ATG ACC GGC TGC GGC AAC ACC CCC ATC TTC AAG GAC GGC CGT GGC TGC 291
Met Thr Gly Cys Gly Asn Thr Thr Pro Ile Phe Lys Asp Gly Arg Gly Cys
55 60 65

GGC TCC TGC TTC GAG ATC AAG TGC ACC AAG CCC GAG TCC TGC TCC GGC 339
Gly Ser Cys Phe Glu Ile Lys Cys Thr Lys Pro Glu Ser Cys Ser Gly
70 75 80 85

```

Fig. 1

GAG GCT GTC ACC GTC ACA ATC ACC GAC GAC AAC GAG GAG CCC ATC GCA	387
Glu Ala Val Thr Val Thr Val Thr Val Thr Val Thr Val Thr Val Thr	100
CCC TAC CAT TTC GAC CTC TCG GGC CAC GCG TTC GGG TCC ATG GCG AAG	435
Pro Tyr His Phe Asp Leu Ser Gly His Ala Phe Gly Ser Met Ala Lys	115
AAG GGC GAG GAG CAG AAG CTC CGC AGC GCC GGC GAG CTG GAG CTC CAG	483
Lys Gly Glu Glu Gln Lys Leu Arg Ser Ala Gly Glu Leu Glu Leu Gln	130
TTC AGG CGG GTC AAG TGC AAG TAC CCG GAC GGC ACC AAG CCG ACA TTC	531
Phe Arg Arg Val Lys Cys Lys Tyr Pro Asp Gly Thr Lys Pro Thr Phe	145
CAC GTC GAG AAG GCT TCC AAC CCC AAC TAC CTC CTC GCT ATT CTG GTG AAG	579
His Val Glu Lys Ala Ser Asn Pro Asn Tyr Leu Ala Ile Leu Val Lys	165
TAC GTC GAC GGC GAC GGT GAC GTG GTG GCG GTG GAC ATC AAG GAG AAG	627
Tyr Val Asp Gly Asp Gly Asp Val Val Ala Val Asp Ile Lys Glu Lys	180

Fig. 1 cont.

GGC AAG GAT AAG TGG ATC GAG CTC AAG GAG TCG TGG GGA GCA GTC TGG	675
Gly Lys Asp Lys Trp Ile Glu Leu Lys Glu Ser Trp Gly Ala Val Trp	195
185	
AGG ATC GAC ACC CCC GAT AAG CTG ACG GGC CCA TTC ACC GTC CGC TAC	723
Arg Ile Asp Thr Pro Asp Lys Leu Thr Gly Pro Phe Thr Val Arg Tyr	210
200	
ACC ACC GAG GGC ACC AAA TCC GAA GTC GAG GAT GTC ATC CCT GAG	771
Thr Thr Glu Gly Gly Thr Lys Ser Glu Val Glu Asp Val Ile Pro Glu	195
215	
GGC TGG AAG GCC GAC ACC TCC TAC TCG GCC AAG TGAGCA	810
Gly Trp Lys Ala Asp Thr Ser Tyr Ser Ala Lys	240
230	
	235

Fig. 1 cont.

PEPTIDE NAME	PEPTIDE SEQUENCE
LPI-1	IAKVPPGPNITA EYGDKWLD
LPI-1.1	IAKVXPGXNITA EYGDKWLD
LPI-2	TA EYGDKWLD AKSTWYGKPT
LPI-3	AKSTWYGKPTGAGPKDNGGA
LPI-4	GAGPKDNGGACGYKNVDKAP
LPI-4.1	GAGPKDNGGACGYKDVDKAP
LPI-5	CGYKDVDKAPFNGMTGCGNT
LPI-6	FNGMTGCGNTPIFKDGRGCG
LPI-7	PIFKDGRGCGSCFEIKCTKP
LPI-8	SCFEIKCTKPESCSGEAVTV
LPI-9	ESCSGEAVTVTITDDNEEPI
LPI-10	TITDDNEEPIAPYHFDLSGH
LPI-11	APYHFDLSGHAFGSMADDGE
LPI-11.1	APYHFDLSGHAFGSMAKKGE
LPI-12	AFGSMADDGEEQKLRSAGEL
LPI-12.1	AFGSMAKKGEEQKLRSAGEL
LPI-13	EQKLRSAGELELQFRRVKCK
LPI-14	ELQFRRVKCKYPDDTKPTFH
LPI-15	YPDDTKPTFHVEKASNPNYL
LPI-16	VEKASNPNYLAILVKYVDGD
LPI-16.1	VEKGSNPNYLAILVKYVDGD
LPI-17	AILVKYVDGDGDVVAVDIKE
LPI-18	GDVVAVDIKEKGKDKWIELK
LPI-19	KGKDKWIELKESWGAVWRID
LPI-20	ESWGAVWRIDTPDKLTGPFT
LPI-21	TPDKLTGPFTVRYTTEGGTK
LPI-22	VRYTTEGGTKSEVEDVIPEG
LPI-23	SEVEDVIPEGWKADTSYSAK

Fig. 2




Fig. 3

PEPTIDE NAME	PEPTIDE SEQUENCE
LPI-16.1	VEKGSNPNYLAILVKYVDGD
LPI-16.2	DEVEKGSNPNYLAILVKYVDGD
LPI-16.3	DEAEKGSNPNYLAILVKYVDGD
LPI-16.4	KKVEKGSNPNYLAILVKK
LPI-16.5	VEKGSNPNYLAILDE
LPI-16.6	AEKGSNPNYLAILDE
LPI-16.7	DEVEKGSNPNYLAIDE
LPI-16.9	KKAEEKSNPNYLAILVKK
LPI-16.10	DEPNYLAILVKYVDE
LPI-18	GDVVAVDIKEKGKDKWIELK
LPI-18.5	GDVVAVDIKEKGKDK
LPI-18.6	VAVDIKEKGKDKWIE
LPI-18.7	AVDIKEKGKDKWIEL
LPI-18.8	DIKEKGKDKWIELK
LPI-20	ESWGAVWRIDTPDKLTGPFT
LPI-20.2	WGAVWRIDTPDKLT
LPI-20.3	GAVWRIDTPDKLTG
LPI-20.4	WRIDTPDKLTGPFT
LPI-20.5	ESWGAVWRIDTPDK
LPI-20.6	AGAVWRIDTPDKLT
LPI-23	SEVEDVIPEGWKADTSYSAK
LPI-23.1	SEVEDVIPEGWKADT
LPI-23.2	EDVIPEGWKADTSYS
LPI-23.4	IPEGWKADTSYSAK

Fig. 4

ATCCCCGAAGGTGCCCCCGGGCCCGAACAATCACGGCGACCTACGGTGACAAAGTGGCTGGAC 60
I P K V P P P G P N I T A T Y G D K W L D
1 10 20

GCGAAGAGCACATGGTACGGCAAGCCGACGGCGCCCGGCCCAAGGACAACGGCGGCGCG 120
A K S T W Y G K P T G A G P K D N G G A
30 40

TGCGGGTACAAGGACGTGGACAAGGCGCGTTCACGGCATGACCGGGTGCGGCAACACC 180
C G Y K D V D K A P F N G M T G C G N T
50 60

CCCATCTTCAAGGACGGCGGGGTGCGGTTCCTGCTTCGAGATCAAGTGCACGAAGCCC 240
P I F K D G R G C G S C F E I K C T K P
70 80

GAGTCGTGCTCCGGCGAGGCCGTCAACCGTCCACATCACCGACGACAACGAGGCCCATC 300
E S C S G E A V T V H I T D D N E E P I
90 100

GCGCCCTACCACTTCGACCTTTCGGGCCACGCGTTCGGTTCATGGCGAAGAGGCGGAG 360
A P Y H F D L S G H A F G S M A K K G E
110 120

GAGCAGAAGCTGCGCAGCGCGGGCGAGCTGGAGCTGCAGTTTAGCGGGGTGAAGTGCAAG 420
E Q K L R S A G E L E L Q F R R V K C K
130 140

Fig. 5

TACCCCGAGGCACCAAGGTGACCTTCCACGTCGAGAAGGGTTCCAAACCCCAACTACCTG 480
Y P E G T K V T F H V E K G S N P N Y L 160
150

GGCTGCTGGTGAAGTACGTCGACGGCGACGGCGACGTCGTCGGTGGCGGTGATATCAAGGAG 540
A L L V K Y V D G D G D V V A V D I K E 180
170

AAGGCAAGGACAAGTGGATCGCGCTCAAGGAGTCATGGGGAGCCATCTGGAGGGTGGAC 600
K G K D K W I A L K E S W G A I W R V D 200
190

ACCCCGACAAGCTGACGGGCCCATTCACCGTTCGCTACACCCAGGAGGCACCAAG 660
T P D K L T G P F T V R Y T T E G G T K 220
210

TCCGAAGTTGAGGACGTCATCCCCGAGGGCTGGAAGGCCGACGCTACGAGTCCAAG 720
S E V E D V I P E G W K A D A S Y E S K 240
230

TGA 723

Fig. 5 cont.

ATCGGAAGGTTCCCGGCCCGAACAATCACGGGACCTACGGCGACAAGTGGCTTGAC 60
I A K V P P G P N I T A T Y G D K W L D 20
1 10

CGGAAGACACCTGGTACGGCAAGCCGACCGGNBCCGGTCCCAAGGACAACGGCGCGG 120
A K S T W Y G K P T G X G P K D N G G A 40
30

TGCGGATACAAGGACGTGGACAAGCCCCCGTTACGGGGCATGACCGGCTGCGGCAACACC 180
C G Y K D V D K P P F S G M T G C G N T 60
50

CCCATCTTCAAGTCCGGCGCGGCTGCGGCTCCTGTTCGAGATCAAGTGACCAAGCCC 240
P I F K S G R G C G S C F E I K C T K P 80
70

GAGTCCTGCTCCGGGGAGCCCGTCCGTCCACATCACCGACGACAACGAGGAGCCCATC 300
E S C S G E P V L V H I T D D N E E P I 100
90

GCCGCCCTACCACTTCGACCTCTCCGGCAAGGCGTTCGGGGCCATGGCCAAGAGGTGAG 360
A A Y H F D L S G K A F G A M A K K G E 120
110

Fig. 6

GAGCAGAAAGCTGCGCAGCGCGCGGAGCTGGAGCTCAAGTTCCGCCGCGTCAAGTGCGAG 420
 E Q K L R S A G E L E L K F R R V K C E 140
 130
 TACCCGAAGGGACCAAGGTTACCTTCCACGTCGAGAGGGGTCCAACCCCACTACCTT 480
 Y P K G T K V T F H V E K G S N P N Y L 160
 150
 GCGCTGCTGGTGAAGTACGTCGACGGCGGACGGGACGTCGTGGCGGTGGACATCAAGCAG 540
 A L L V K Y V D G D G D V V A V D I K Q 180
 170
 AAGGCAAGGACAAAGTGGATCGAGCTCAAGGAGTCGTGGGAGCCGTCCTGGAGGATCGAC 600
 K G K D K W I E L K E S W G A V W R I D 200
 190
 ACCCCGACAAGCTACCGGCCCCCTTACCGTCGCTACACCCGAGGGCGGCCAACAG 660
 T P D K L T G P F T V R Y T T E G G T K 220
 210
 GCCGAAGCCGAGGACGTATCCCCGAGGGCTGGAAGGCCGACACCGCCTACGAGGCCAAG 720
 A E A E D V I P E G W K A D T A Y E A K 240
 230
 TGA 723

Fig. 6 cont.

ATCGCGAAGGTGCCCCGGGTCCGAACATCACGGCGACCTACGGCGACAAGTGGCTCGAC 60
I A K V P P G P N I T A T Y G D K W L D 20
1 10
GCGAAGAGCACATGGTACGGCAAGCCGACGGGGCCGGTCCCAAGGACAACGGCGGCGCT 120
A K S T W Y G K P T G A G P K D N G G A 40
30
TGC GGGTACAAGGACGTGGACAAGCCCCCGTTACGGGGCATGACCGGCTGCGGCAACACC 180
C G Y K D V D K P P F S G M T G C G N T 60
50
CCCATCTTCAAGTCCGGCCGTGGCTGCGGCTCCTGTCTTGAGATCAAGTGCACGAAGCCC 240
P I F K S G R G C G S C F E I K C T K P 80
70
GAGGCCTGCTCCGGCGAGCCCGTGGTAGTCCACATCACCGACGACAACGAGAGCCCATC 300
E A C S G E P V V V H I T D D N E E P I 100
90
GCCCCCTACCAC'TTCGACCTCTCCGGCCACGCGTTCGGGGCGATGGCCAAAGAAGGCGGAT 360
A P Y H F D L S G H A F G A M A K K G D 120
110

Fig. 7

GAGCAGAAGCTGCGCACGGCCGGCGAGCTGGAGCTCCAGTTCGGCGCGTCAAGTCAAG 420
E Q K L R T A G E L E L Q F R R V K C K 140
130
TACCGGAGGGACCAAGGTGACCTTCCACGTGGAGAAGGGGTCCAACCCCACTACCTG 480
Y P E G T K V T F H V E K G S N P N Y L 160
150
GCGCTGCTTGTGAAGTACGTTAACGGCGACGGAGACGTGGTGGCGGTGGACATCAAGGAG 540
A L L V K Y V N G D G D V V A V D I K E 180
170
AAGGCAAGGACAAGTGGATCGAGCTCAAGGAGTCGTGGGGAGCCATCTGGAGGATCGAC 600
K G K D K W I E L K E S W G A I W R I D 200
190
ACTCCCGACAAGCTCAGGGCCCTTCAACCGTCCGCTACACCAGGCGGCACCAAG 660
T P D K L T G P F T V R Y T T E G G T K 220
210
ACCGAAGCCGAGGACGTCAATCCCTGAGGGCTGGAAGGCCGACACCAAGTACGAGTCCAAG 720
T E A E D V I P E G W K A D T S Y E S K 240
230
TGA 723

Fig. 7 cont.

1	10	20	30	40	50	60	
IAKVP	PGPNITAEY	GDKWLD	AKSTWYGKPTG	AGPKDNG	GACGYK ^D	VDKAPFNGMTGCGNT	Lol p I
-P-	T-	T-	T-	T-	T-	T-	Dac g I
-	-	-	-	-	-	-	Phl p I
-	-	-	-	-	-	-	Poa p I
70	80	90	100	110	120		
PIFKD	GRGSGCFEIK	CTKPESCSGE	AVT ^D VTITDD	NEEPIAPYHFD	LSGHAFGSM	AKKGE	Lol p I
-	-	-	-	-	-	-	Dac g I
-	-	-	-	-	-	-	Phl p I
-	-	-	-	-	-	-	Poa p I
130	140	150	160	170	180		
EQKLR	SAGELELQFRR	VCKY ^D PD ^G	TKPTFHVEK ^C	SNPNYLAILV	KYVDGDV	VAVDIKE	Lol p I
-	-	-	-	-	-	-	Dac g I
-	-	-	-	-	-	-	Phl p I
-	-	-	-	-	-	-	Poa p I
190	200	210	220	230	240		
KGKDKW ^I	ELKESGAVWR	IDTPDKLTG	PFTVRYTTEG	TKSE ^V	VEDVIPEGW	KADTSYSAK	Lol p I
-	-	-	-	-	-	-	Dac g I
-	-	-	-	-	-	-	Phl p I
-	-	-	-	-	-	-	Poa p I

Fig. 8

PEPTIDE NAME	PEPTIDE SEQUENCE
LPI-16.1	VEKGSNPNYLAILVKYVDGD
LPI-16.8	VEKGSNPNYLAILVKYVDGD
LPI-16.N	VEKGSNPNYLAILVKYVNGD
LPI-18	GDVVAVDIKEKGDKKWIELK
LPI-18.9	GDVVAVDIKEKGDKKWIALK
LPI-18.X	GDVVAVDIKQKGDKKWIELK
LPI-20	ESWGAVWRIDTPDKLTGPFT
LPI-20.7	ESWGAIWRIDTPDKLTGPFT
LPI-20.Y	ESWGAIWRVDTPDKLTGPFT
LPI-23	SEVEDVIPEGWKADTSYSAK
LPI-23.5	TEAEDVIPEGWKADTSYESK
LPI-23.6	AEAEDVIPEGWKADTAYEAK
LPI-23.Z	SEVEDVIPEGWKADASYESK
LPI-23.ZZ	SEVEDVIPEGWKADTSYESK

Fig. 9